

## Chemical constituents and analgesic activity of *Telfaria occidentalis*

Jude Efiom Okokon<sup>1</sup>, Ashana Dar<sup>2</sup>, Muhammad Iqbal Choudhary<sup>2</sup>

<sup>2</sup>Department of Pharmacology and Toxicology, Faculty of Pharmacy, University of Uyo, Uyo, Nigeria.

<sup>2</sup>International Centre for Chemical and Biological Sciences, University of Karachi, Karachi-75270, Pakistan.

\*Corresponding author: judeefiom@yahoo.com; Tel: +234-802-3453678

**Received:** 4 July 2012, **Revised:** 18 July 2012, **Accepted:** 19 July 2012

### Abstract

The seed of *Telfairia occidentalis* is use commonly for nutritional purposes and medicinally in the treatment of malaria and inflammatory diseases. The ethanolic seed extract of *Telfairia occidentalis* (450 -1350 mg/kg) was evaluated for analgesic activity using acetic acid-induced writhing, formalin-induced hind paw licking and thermal –induced pain models. The hexane and dichloromethane fractions were also analyzed using Gas chromatography-mass spectrometry (GCMS). The seed extract exhibited a dose-dependent inhibition of pains in the three experimental models. The GCMS analysis revealed the presence of pharmacological active compounds which are responsible for the analgesic activity.

**Keywords:** *Telfairia occidentalis*; analgesic; vegetable; GCMS; Chemical constituents

### Introduction

*Telfairia occidentalis* (Hook. F) Vahl. popularly known as fluted pumpkin is a member of Cucurbitaceae family. The plant is cultivated in Southern Nigeria mainly for the leaves and seeds which are eaten because of their high content of protein, vitamins and minerals (Johnson and Johnson, 1996). *T. occidentalis* leaf are often used as vegetable in the preparation of soups, while the seeds are eaten raw or roasted and also ground into powder and used as soup thickening. Reports of hypoglycaemic and antidiabetic activities (Aderibigbe et al., 1999; Alada, 2000; Eseyin et al., 2000; Eseyin et al., 2005; Nwozo et al., 2004), antioxidant and antimicrobial activities (Obboh et al., 2006) of the leaf have been published. Several workers have reported on the nutritional composition, chemical characterization and functional properties of fluted pumpkin seed (Asiegbu, 1987; Badifu et al., 1995; Agatemor, 2006; Ezugwu and Nwodo, 2000; Fagbemi et al., 2005). The seed has been reported to possess antiplasmodial (Okokon et al., 2008). Most researches have focused on the leaf and information on the medicinal properties of the seed is scanty. We report in this study the analgesic activity as well as GC-MS analysis of hexane and dichloromethane fractions of the seed extract of *Telfairia occidentalis* from Nigeria.

## Materials and methods

### *Plants collection*

The plant material *Telfairia occidentalis* (seeds) were bought from local markets in Uruan area, Akwa Ibom State, Nigeria in April, 2011. The plant was identified and authenticated by Dr. Magaret Bassey of Department of Botany and Ecological Studies, University of Uyo, Uyo, Nigeria. A voucher specimen was deposited in the faculty of Pharmacy Herbarium, University of Uyo, Uyo with voucher no. FPHUU 110.

### *Extraction*

The seeds were washed and shade-dried for two weeks. The dried plants' materials were further chopped into small pieces and reduced to powder. The powdered material was macerated in 70 % ethanol. The liquid filtrates were concentrated and evaporated to dryness in vacuo 40°C using rotary evaporator. The crude ethanolic extract (100 g) was further partitioned successively into 1L each of n-hexane, dichloromethane, ethyl acetate and butanol to give the corresponding fractions of these solvents.

### *Evaluation of analgesic activity*

#### *Acetic acid induced writhing in mice*

The abdominal constrictions resulting from intraperitoneal (i.p) injection of 3% acetic acid consisting of the contraction of abdominal muscles together with the stretching of hindlimbs, was carried out according to the procedure described by Santos et al. (1994), Correa et al. (1996) and Nwafor et al.,(2010). The animals were divided into 5 groups of 6 mice per group. Group 1 served as negative control and received 10ml/kg of normal saline, while groups 2, 3 and 4 were pre-treated with 450, 900 and 1350 mg/kg doses of *Telfairia occidentalis* seed extract intraperitoneally, and group 5 received 100 mg/kg of acetyl salicylic acid. After 30 minutes, 0.2 ml of 2% acetic acid was administered intraperitoneally (i.p). The number of writhing movements was counted for 30 minutes. Antinociception (analgesia) was expressed as the reduction of the number of abdominal constrictions between control animals and mice pretreated with extracts.

#### *Formalin-induced hind paw licking in mice*

The procedure was essentially similar to that described by Hunskaar and Hole (1987), Correa and Calixto (1993), Gorki et al., (1993) and Okokon and Nwafor,(2010). The animals were used to analyze the first phase of formalin-induced licking and 20µL of 2.5% formalin solution (0.9% formaldehyde) made up in phosphate buffer solution (PBS concentration: NaCl 137mM, KCl 2.7mM and phosphate buffer, 10 mM) was injected subcutaneously under the surface of the right hind paw. The amount of time spent licking the injected paw was timed and considered as indication of pain. The first of the nociceptive response normally peaks 5 minutes after injection and the second phase 15 - 30 minutes after formalin injection, representing the neurogenic and inflammatory pain responses, respectively (Hunskaar and Hole, 1987). Adult albino mice (23 – 27 g) of either sex randomised into five groups of 6 mi-

ce each were used for the experiment. The mice were fasted for 24 hours before used but allowed access to water. The animals in group 1 (negative control) received 10ml/kg of normal saline, groups 2 - 4 received 450, 900 and 1350 mg/kg doses of the seed extract, while group 5 received 100mg/kg of acetyl salicylic acid 30 minutes before being challenged with buffered formalin. The responses were measured for 5mins after formalin injection (first phase) and 15–30 mins after formalin injection (second phase).

#### *Thermally induced pain in mice*

The effect of extract on hot plate induced pain was investigated in adult mice. The hot plate was used to measure the response latencies according to the method of Vaz et al., (1996) and Okokon and Nwafor, (2010). In these experiments, the hot plate was maintained at  $45\pm 1^{\circ}\text{C}$ , each animal was placed into a glass beaker of 50 cm diameter on the heated surface, and the time(s) between placement and shaking or licking of the paws or jumping was recorded as the index of response latency. An automatic 30-second cut-off was used to prevent tissue damage. The animals were randomly divided into 5 groups of 6 mice each and fasted for 24 hours but allowed access to water. Group 1 animal served as negative control and received 10ml/kg of normal saline. Groups 2, 3 and 4 were pre-treated intraperitoneally with 450, 900 and 1350 mg/kg doses of *Telfairia occidentalis* seed extract respectively, while group 5 animals received 100 mg/kg of acetyl salicylic acid intraperitoneally, 30 min-utes prior to the placement on the hot plate.

#### *GC-MS analysis of hexane and dichloromethane fraction*

Quantitative and qualitative data were determined by GC and GC-MS, respectively. Each fraction was injected onto a Shimadzu GC-17A system, equipped with an AOC-20i autosampler and a split/ splitless injector. The column used was an DB-5 (Optima-5), 30 m, 0.25 mm i.d., 0.25  $\mu\text{m}$  df, coated with 5 % diphenyl-95 % polydimethylsiloxane, operated with the following oven temperature programme:  $50^{\circ}\text{C}$ , held for 1 min, rising at  $3^{\circ}\text{C}/\text{min}$  to  $250^{\circ}\text{C}$ , held for 5 min, rising at  $2^{\circ}\text{C}/\text{min}$  to  $280^{\circ}\text{C}$ , held for 3 min; injection temperature and volume,  $250^{\circ}\text{C}$  and 1.0  $\mu\text{l}$ , respectively; injection mode, split; split ratio, 30:1; carrier gas, nitrogen at 30 cm/s linear velocity and inlet pressure 99.8 KPa; detector temperature,  $280^{\circ}\text{C}$ ; hydrogen, flow rate, 50 ml/min; air flow rate, 400 ml/min; make-up ( $\text{H}_2/\text{air}$ ), flow rate, 50 ml/min; sampling rate, 40 ms. Data were acquired by means of GC solution software (Shimadzu). Agilent 6890N GC was interfaced with a VG Analytical 70-250 s double -focusing mass spectrometer. Helium was used as the carrier gas. The MS operating conditions were: ionization voltage 70 eV, ion source  $250^{\circ}\text{C}$ . The GC was fitted with a 30 m x 0.32 mm fused capillary silica column coated with DB-5. The GC operating parameters were identical with those of GC analysis described above.

The identification of components present in the various active fractions of the plants' extracts was based on direct comparison of the retention times and mass spectral data with those for standard compounds, and by computer matching with the Wiley 229 and Nist 21 Library, as well as by comparison of the fragmentation patterns of the mass spectra with those reported in the literatures (Adams, 2001; Setzer et al., 2007).

## Statistical analysis

Data obtained from this work were analyzed statistically using Students' t-test and ANOVA (One- or Two-way) followed by a post test (Tukey-Kramer multiple comparison test). Differences between means will be considered significant at 1% and 5% level of significance i.e  $P < 0.01$  and  $0.05$ .

## Results

### Effect of on acetic acid-induced writhing in mice

The extract (450–1350 mg/kg) demonstrated a dose-dependent reduction in acetic acid-induced writhing in mice. The reductions were statistically significant ( $p < 0.001$ ) relative to control and comparable to that of the standard drug, ASA, at the highest dose, 1350 mg/kg. (Table 1).

### Effect on formalin-induced hind paw licking in mice

The extract exhibited a dose-dependent effect on formalin-induced hind paw licking in mice. This inhibition was significant relative to the control ( $p < 0.001$ ) and comparable to that of the standard drug, ASA, at the highest dose, 1350 mg/kg. (Table 2).

### Effect on thermally-induced pain in mice

The extract exhibited a dose-dependent effect on thermally-induced pain in mice. This inhibition was statistically significant ( $p < 0.001$ ) relative to the control (Table 3).

Table 1. Effect of *Telfairia occidentalis* on acetic acid induced writhing in mice.

Dose (mg/kg)	Time interval (hr)						
	5	10	15	20	25	30	Total
Control	5.00 ± 1.06	8.83 ± 1.42	19.16 ± 2.13	18.16 ± 1.49	15.7 ± 0.71	12.33 ± 1.20	79.14 ± 3.03
Extract 450	0.00 <sup>c</sup>	9.60 ± 0.88	11.16 ± 2.27 <sup>a</sup>	12.50 ± 1.36 <sup>a</sup>	11.5 ± 0.23	10.66 ± 1.82	55.42 ± 3.26 <sup>c</sup>
Extract 900	0.00 <sup>c</sup>	9.16 ± 1.40	11.0 ± 0.96 <sup>a</sup>	11.16 ± 0.94 <sup>a</sup>	8.50 ± 1.23 <sup>a</sup>	6.80 ± 1.37 <sup>a</sup>	46.62 ± 2.23 <sup>c</sup>
Extract 1350	0.00 <sup>c</sup>	9.50 ± 1.54	11.08 ± 0.67 <sup>a</sup>	9.50 ± 1.28 <sup>b</sup>	7.83 ± 1.53 <sup>a</sup>	3.83 ± 0.87 <sup>b</sup>	41.74 ± 2.33 <sup>c</sup>
ASA 100	0.00 <sup>c</sup>	2.00 ± 0.00 <sup>b</sup>	6.40 ± 1.93 <sup>c</sup>	9.50 ± 1.73 <sup>b</sup>	9.50 ± 1.28 <sup>b</sup>	9.30 ± 1.85 <sup>b</sup>	36.70 ± 2.56 <sup>c</sup>

Data are expressed as mean ± SEM. Significant at <sup>a</sup> $P < 0.05$ , <sup>b</sup> $P < 0.001$  when compared to control. n = 6.

Table 2. Effect of *Telfairia occidentalis* on formalin –induced hind paw licking in mice.

Dose (mg/kg)	Time interval (hr)						
	5	10	15	20	25	30	Total
Control	35.46 ± 0.19	15.27 ± 0.60	16.10 ± 0.24	12.44 ± 0.54	8.35 ± 0.38	5.88 ± 0.11	87.94 ± 3.25
Extract 450	29.50 ± 0.88	6.00 ± 0.25 <sup>b</sup>	5.00 ± 0.36 <sup>c</sup>	4.33 ± 0.42 <sup>c</sup>	4.00 ± 0.36 <sup>c</sup>	3.66 ± 0.21 <sup>c</sup>	52.49 ± 2.71 <sup>c</sup>
Extract 900	24.16 ± 0.28 <sup>c</sup>	5.33 ± 0.21 <sup>c</sup>	4.83 ± 0.33 <sup>c</sup>	4.16 ± 0.16 <sup>c</sup>	3.33 ± 0.21 <sup>c</sup>	3.00 ± 0.36 <sup>c</sup>	44.76 ± 1.61 <sup>c</sup>
Extract 1350	18.5 ± 2.74 <sup>c</sup>	0.00 <sup>c</sup>	0.50 ± 0.49 <sup>c</sup>	3.00 ± 0.516 <sup>c</sup>	2.50 ± 0.22 <sup>c</sup>	2.50 ± 0.22 <sup>c</sup>	27.00 ± 4.16 <sup>c</sup>
ASA 100	8.83 ± 0.43 <sup>c</sup>	1.66 ± 0.21 <sup>c</sup>	2.80 ± 0.16 <sup>c</sup>	2.16 ± 0.16 <sup>c</sup>	1.16 ± 0.16 <sup>c</sup>	0.00 <sup>c</sup>	16.61 ± 0.43 <sup>c</sup>

Data are expressed as mean ± SEM. Significant at <sup>a</sup> $P < 0.05$ , <sup>b</sup> $P < 0.001$  when compared to control. n = 6.

Table 3. Effect of *Telfairia occidentalis* on hot-plate test in mice.

Group	Dose (Mg/kg)	Reaction time (sec)(Mean±SEM)	% inhibition
Control	-	5.10±0.29	-
<i>T. occidentalis</i>	450	6.95±0.81 <sup>a</sup>	36.27
<i>T. occidentalis</i>	900	10.77±0.75 <sup>a</sup>	111.17
<i>T. occidentalis</i>	1350	16.38±0.40 <sup>b</sup>	221.17
ASA	100	18.04±0.25 <sup>b</sup>	253.72

Data are expressed as mean±SEM. Significant at <sup>a</sup>P < 0.05, <sup>b</sup>P < 0.001 when compared to control. n = 6.

### GC-MS analysis

The GCMS analysis of the hexane and dichloromethane fractions of *Telfairia occidentalis* seed revealed the presence of 11 bioactive compounds each as represented in Tables 4 and 5.

### Discussion

The seed of *Telfairia occidentalis* use by the Ibibios of Niger Delta of Nigeria for nutritive purposes in the making of soup was evaluated for antinociceptive activity in rodent. The seed extract in this study, significantly reduced acetic acid-induced writhing, formalin-induced hind paw licking as well as delayed the reaction time of animals (mice) to thermally induced pain. Acetic acid causes inflammatory pain by inducing capillary permeability (Am-

Table 4. GC-MS analysis of dichloromethane fraction of *Telfairia occidentalis* seeds.

S/No.	Compound	Mol. Wt	Chemical Formula	RI
1.	Pentadecanoic acid, 14-methyl-,methyl ester	270	C <sub>17</sub> H <sub>34</sub> O <sub>2</sub>	619
2.	Hexadecanoic acid	256	C <sub>16</sub> H <sub>32</sub> O <sub>2</sub>	648
3.	8,11-Octadecadienoic acid, methyl ester	294	C <sub>19</sub> H <sub>34</sub> O <sub>2</sub>	703
4.	16-Octadecenoic acid,methyl ester	296	C <sub>19</sub> H <sub>36</sub> O <sub>2</sub>	706
5.	Heptadecanoic acid,6-methyl, methyl ester	298	C <sub>19</sub> H <sub>38</sub> O <sub>2</sub>	719
6.	9, 12-Octadecadienoyl chloride(Z,Z)-	298	C <sub>18</sub> H <sub>31</sub> ClO	735
7.	9-Octadecadienoic acid (Z)-,2,3-dihydroxypropyl ester	356	C <sub>21</sub> H <sub>40</sub> O <sub>4</sub>	741
8.	Octadecanoic acid	284	C <sub>18</sub> H <sub>36</sub> O <sub>2</sub>	749
9.	Hexadecanoic acid, 2,3-bis[(trimethylsilyl)oxy] propyl ester	474	C <sub>25</sub> H <sub>54</sub> O <sub>4</sub> Si <sub>2</sub>	876
10.	Cyclohexanespiro-5'-(4'-methyl-2'-phenyl-2'-oxazoline	229	C <sub>15</sub> H <sub>19</sub> NO	947
11.	9-Octadecenoic acid (Z)-,2-hydroxy-1-(hydroxymethyl) ethyl ester.	356	C <sub>21</sub> H <sub>40</sub> O <sub>4</sub>	966

Table 4. GC-MS analysis of n-hexane fraction of *Telfairia occidentalis* seeds.

S/No.	Compound	Mol. Wt	Chemical Formula	RI
1.	2,4-Heptadien-6-ynal,(E,E)-	106	C <sub>7</sub> H <sub>6</sub> O	190
2.	Benzoic acid	122	C <sub>7</sub> H <sub>6</sub> O <sub>2</sub>	195
3.	Dodecanoic acid	200	C <sub>12</sub> H <sub>24</sub> O <sub>2</sub>	201
4.	Linoleic acid ethyl ester	308	C <sub>20</sub> H <sub>36</sub> O <sub>2</sub>	729
5.	Hexadecanoic acid, methyl ester	284	C <sub>18</sub> H <sub>36</sub> O <sub>2</sub>	756
6.	α-phellandrene	136	C <sub>10</sub> H <sub>16</sub>	1005
7.	α-campholene aldehyde	152	C <sub>19</sub> H <sub>16</sub> O	1123
8.	Terpinen-4-ol	154	C <sub>10</sub> H <sub>18</sub> O	1137
9.	Trans-β-ocimene	136	C <sub>10</sub> H <sub>16</sub>	1150
10.	Borneol	154	C <sub>10</sub> H <sub>18</sub> O	1164
11.	Stigmastan-3-ol, 5-chloro-,acetate,(3a',5a')-	492	C <sub>13</sub> H <sub>53</sub> ClO <sub>2</sub>	1175

Amico-Roxas *et al.*, 1984; Nwafor *et al.*, 2007), and in part through local peritoneal receptors from peritoneal fluid concentration of PGE<sub>2</sub> and PGF<sub>2</sub>α (Deraedt *et al.*, 1980; Bentley *et al.*, 1983). This model is used to distinguish between central and peripheral pain.

Formalin exhibits neurogenic and inflammatory pains (Vaz *et al.*, 1996, 1997) and measures both centrally and peripherally mediated activities that are characteristic of biphasic pain response. The first phase of formalin-induced hind paw licking is selective for centrally acting analgesics such as morphine (Berken *et al.*, 1991), while the late phase of formalin-induced hind paw licking is peripherally mediated. Analgesic (nociceptive) receptors mediate both the neurogenic and non-neurogenic pains (Lembeck and Holzer, 1979).

The study also shows that the extract significantly delayed the reaction time of thermally-induced (hot plate) test. This model is selective for centrally acting analgesics and indicates narcotic involvement (Turner, 1995) with opioid receptors.

The GCMS analysis of the hexane fraction has revealed the presence of a-phellandrene, an acyclic monoterpene which has been reported for significant anti-inflammatory and antinociceptive activities against acetic acid-induced writhing and capsaicin test. It is also reported to inhibit both phases of formalin-induced paw licking and in addition decreased carrageenan-induced hyperalgesia (Lima *et al.*, 2012). Moreover, Terpinen-4-ol which has been found in the seed extract has been reported to suppress production of prostaglandin and *in vitro* of TNF-α, IL-1β, as well as IL-8, IL-10 and PGE<sub>2</sub> by LPS-activated human blood monocytes (Hart *et al.*, 2000; Miguel, 2010). These compounds may in part be responsible for the observed analgesic activity of the seed extract.

The antinociceptive activities exerted by this extract may be attributed to the presence of phytochemical compounds as revealed by the GCMS analysis of the seed extract. That the extract inhibited neurogenic and non-neurogenic pains as well as narcotic pains may in part explain the mechanisms of its action and these effects are due to the presence of phytochemical components in the extract. In conclusion, the seed extract has analgesic property which is due to its phytochemical components. Hence, can be exploited in the management of pains.

### Acknowledgement

The authors are grateful to TWAS for provision of financial for postdoctoral studies and ICCBS, Pakistan for the provision of research facilities.

### Conflict of interest

There is no conflict of interest associated with the authors of this paper.

### References

- Akah PA, Nwanbie A. (1994). Evaluation of Nigerian traditional medicines plants used for rheumatic (inflammatory) disorder. *Journal of Ethnopharmacology* 42, 179 – 182.
- Amico-Roxas M, Caruso A, Trombadore S, Scifo R, Scapagnime U. (1984). Gangliosides antinociceptive effects in rodents. *Archives of International Pharmacodynamics and Therapeutics*. 272, 103-117.

- Bentley GA, Newton SH, Starr J. (1983). Studies on the antinociceptive action of agonist drugs and their interaction with opioid mechanisms. *British Journal of Pharmacy* 79, 125 - 134.
- Berken T, Ostunes L, Lermioglu F, Ozer A. (1991). Antiinflammatory analgesic and antipyretic effect of an aqueous extract of *Erythraea ceulaurum*. *Planta Medica*. 57, 34 -37.
- Besra SE, Sharma RM, Gomes A. (1996). Antiinflammatory effect of petroleum ether extract of leaves of *Litchi Chinensis*. Caertn (sapindaceae). *Journal of Ethnopharmacology* 54, 1-6.
- Correa CR, Calixto JB. (1993). Evidence of participation of  $\beta_1$  and  $\beta_2$  receptors, in formalin induced nociceptive response in mouse. *British Journal of Pharmacology* 110, 193-198.
- Correa CR, Kyle DJ, Chakravarty S, Calixto JB. (1996). Antinociceptive profile of the pseudopeptide  $\beta_2$  bradykinin receptors antagonist NPC 18688 in mice. *British Journal of Pharmacology*.117, 552-556.
- Deraedt R, Jougney S, Falhout M. (1980). Release of Prostaglandin E and F in an algogenic reaction and its inhibition. *European Journal of Pharmacy*. 51, 17-24.
- Ekpendu TO, Akah PA, Adesomoju AA, Okogun JI. (1994). Antiinflammatory and antimicrobial activities of *Mitracarpus scaber* extracts. *International Journal of Pharmacology*. 32, 191-195.
- Gorski F, Correa CR, Filhe VC, Yunes RA, Calixto JB. (1993). Potent antinociceptive activity of a hydroalcoholic extract from *Phyllanthus corcovadensis*. *Journal of Pharmacy and Pharmacology*. 45, 1046-1049.
- Hess SM, Milonig RC. (1972). Inflammation In: Lepow, L. H., Ward, P.S. (Eds). Inflammation, Mechanism and control. Academic Press, New-york, USA. pp.1-2.
- Hunskar S, Hole K. (1987). The formalin test in mice. Dissociation between inflammatory pain. *Pain* 30, 103-114.
- Lembeck F, Holzer P. (1979). Substance-P as neurogenic mediator of antidromic, vasodilatation and neurogenic plasma extravasation. *Naunyn-Schmiedeberg's Archives of Pharmacy*. 310, 175-183.
- Liang YC, Huang YT, Tsau SH, Lin-Shiau SY, Chen CF, Lin JK. (1999). Suppression of inducible cyclo-oxygenase and inducible nitric acid synthase by apigenin and related flavonoid in mouse macrophages. *Carcinogenesis*. 20, 1945-52.
- Lima DF, Brando MS, Moura JB, Leitao JM, Carvalho FA, Miura LM, Leite JS, Sousa DP, Almeida FR.(2012). Antinociceptive activity of the monoterpene  $\alpha$ -phellandrene in rodents. *Journal of Pharmacy and Pharmacology* 64, 283 -292.
- Lin LL, Lin AY, Knopt JL. (1992). Cytosolic phospholipase  $A_2$  is coupled to hormonally regulated release of arachidonic acid. *Proceeding of National Academy of Science, U. S. A.* 89, 6147-6157.
- Mbagwu HO, Anene RA, Adeyemi OO. (2007). Analgesic, antipyretic and antiinflammatory properties of *Mezoneuron benthamianum* Baill Caesalpiniaceae. *Nigerian Quarterly Journal of Hospital Medicine*. 17, 35-41.
- Nwafor PA, Jacks TW, Ekanem AU.. (2007). Analgesic and anti-inflammatory effects of methanolic extract of *Pausinystalia mecroceras* stem bark in rodents. *Journal of Pharmacology* 3, 86-90.
- Nwafor PA, Nwajiobi N, Uko IE, Obot JS. (2010). Analgesic and anti-Inflammatory activities of an ethanol extract of *Smilax krausiana* leaf in mice. *African Journal Biomedical Research* 13, 141 -148.
- Ojewole, J. A. O.(2005). Antinociceptive, antiinflammatory and antidiabetic effects of *Bryophyllum pinnatum* (Crassulaceae) leaf aqueous extract. *Journal of Ethnopharmacology*. 99, 13-19.
- Okokon JE, Nwafor PA. (2010). Antiinflammatory, analgesic and antipyretic activities of ethanolic root extract of *Croton zambesicus*. *Pakistan Journal of Pharmaceutical Sciences*. 23, 383-390.
- Okokon JE., Ekpo A., Eseyin OA. (2009). Antiplasmodial activities of leaf and seed extracts of *Telfairia occidentalis* on *Plasmodium berghei* infection in mice. *Journal of Medicinal Food*.12, 649 - 653.

- Parmer NS, Ghosh MN. (1978). Anti-inflammatory activity of gossypin a biflavonoid isolated from *Hibiscus vitifolicus* Linn. *Indian Journal of Pharmacology*. 10, 277-293.
- Sanchez-Mateo CC, Bonkanka CX, Hernandez-Perez M, Rabanal R M.(2006). Evaluation of the analgesic and topical antiinflammatory effects of *Hypericum reflexum* L. fil. *Journal of Ethnopharmacology* 107, 1-6.
- Santos AR, Cechinel FV, Niero R, Viana AM, Moreno FN, Campos MM, Yunes RA, Calixto JB. (1994). Analgesic effects of callus culture from selected species of *Phyllanthus*. *Journal of Pharmacy and Pharmacology*. 46, 755-759.
- Tjolsen A, Berge OG, Hunskaar S, Rosland JH, Hole K. (1992). The formalin test: An evaluation of the method. *Pain* 51, 5-17.
- Turner RA. (1995). Screening methods in Pharmacology. Vol 1. Academic Press. New York. pp. 85-106.
- Vane T, Booting R. (1987). Inflammation and Mechanism of action of anti-inflammatory drugs. *FASSEB J.* 1, 89-96.
- Vaz ZR, Cechinel V, Yunes RA, Calixto JB. (1996). Antinociceptive action of 2-(4-bromo-benzoyl)-3-methyl-4-6-dimethoxybezofuran, a novel xanthoxyline derivative of chemical and thermal models of nociception in mice. *Journal of Pharmacy and Experimental Therapeutics* 278, 304 - 312.
- Vaz ZR, Risley EA, Calixto JB. (1997). Analgesic effect of the herbal medicine catuama in thermal and chemical models of nociception in mice. *Phytotherapy Research* 11, 101-106.
- Winter CA, Risley EA, Nuss GW. 1962. Carrageenin-induced oedema in hind paw of the rats as an assay of anti-inflammatory drugs. *Proceedings of the Society for Experimental Biology and Medicine* 111, 544-547.