

## A Review on the Phytochemistry and Pharmacology of Genus *Tephrosia*

Saad Touqeer<sup>1,\*</sup>, Muhammad Asad Saeed<sup>1</sup>, Muhammad Ajaib<sup>2</sup>

<sup>1</sup>Department of Pharmacy, The University of Lahore, Lahore, Pakistan.

<sup>2</sup>Department of Botany, GC University, Lahore, Pakistan.

\* Corresponding author: saadtouqeer@gmail.com; Tel: +92-322-4899048

**Received:** 2 March 2013, **Revised:** 2 May 2013, **Accepted:** 9 May 2013

### Abstract

The plants of genus *Tephrosia* of family *Leguminosae* are widely distributed in many tropical and subtropical countries of the world and have been used in folk medicine for the treatment of large number of diseases. The importance of this genus is similar to that of other therapeutically renowned genera. This review includes the chemical studies on different species mainly the isolation and identification of flavonoids, rotenoids and study of activity of some isolated compounds and also includes different pharmacological activities like antioxidant, antimicrobial, anticancer, antiplasmodial, anti-inflammatory, larvicidal and toxicity studies of extracts and fractions.

**Keywords:** *Tephrosia*; Flavonoids; Rotenoids; Phytochemistry; Pharmacology; Activity.

### Introduction

Plants have been used for the treatment of diseases from centuries. Natural product chemistry, especially phytochemistry, has become a topic of interest for most of the researchers due to the advantages of the plant derived medicinal compounds over the traditional ways of using herbal plants. Ethnopharmacology plays an important role in the discovery of new biologically active compounds. The process usually starts with searching of useful plants from different records to the development of methods for the industrial production of drugs (Rout et al., 2009; Farnsworth et al., 1985; Koehn and Carter, 2005). According to World Health Organization (WHO) more than 80% of the world's population uses plants for the treatment of their diseases (Calixto et al., 1998; Duraipandiyani et al., 2006).

The genus *Tephrosia* belongs to the family *Leguminosae* and subfamily Papilionaceae. There are approximately 400 species included in this genus. The plants in this genus are widely distributed in tropical, sub-tropical and arid regions of the world (Willis, 1973; Al-

Zahrani, 2007). The plants are prostrate or erect herbs or in the form of soft or woody shrubs (Hacker, 1990). The base chromosome number of this genus is  $X=11$  and are placed in the tribe Galegeae of the family *Leguminosae* (Atchison, 1951; Agarwal and Gupta 1983). Many plants from this genus have been used traditionally for the treatment of diseases like rheumatic pains, syphilis, dropsy, stomach ache, diarrhea, asthma, abortifacient, respiratory disorders, laxative, diuretic, and inflammation etc (Qureshi et al., 2010; Dzenda et al., 2007). *Tephrosia purpurea*, an important plant of the genus is used as tonic, laxative, antivenom, antiulcer, antidiarrheal, and in leprosy (Virupanagouda et al., 2011).

The main purpose of this review is to provide a comprehensive and up-to-date knowledge of the pharmacological and phytochemical research work performed on the genus *Tephrosia*. The plants of this genus have a large potential for study of its activities and chemical constituents for important leads.

### Chemical constituents from plants of genus *Tephrosia*

A great variety of plants belonging to genus *Tephrosia* have been studied for their chemical constituents and pharmacological activities. The number of species studied phytochemically are much more than those studied pharmacologically. Different classes of organic compounds have been isolated of which some have been tested for their biological activities and some still unknown for their effect. The main classes of compounds include flavonoids, rotenoids, terpenoids, and sterols. It should be noted that flavonoids are the most abundantly isolated and identified compounds in the genus. Similarly in case of essential oil and fixed oil, we can see that very little work has been done by the scientists. *Tephrosia purpurea*, *Tephrosia toxicaria*, *Tephrosia candida*, *Tephrosia elata*, and *Tephrosia villosa* have been the plants of interest for the scientists. The readers will also find some work on the stereochemistry of compounds. Praecansone A, flavonoid from *Tephrosia pumila* for example, exists in the form of two isomers (Dagne et al., 2012).

Table 1 briefly describes all the chemical work done on genus *Tephrosia*. Readers will see some novel compounds as well some compounds repeating in many species. Some chemicals in the table would be familiar to us as they have been isolated from other genera as well

Table 1. Chemical constituents from plants of genus *Tephrosia*.

Species	Class	Compound	Reference
<i>Tephrosia abbotiae</i>	Flavonoid	abbotin	Gómez-Garibay et al., 1986
<i>Tephrosia aequilata</i>	Flavonoid	tephrobotin obovatin methyl ether (E)-praecansone A demethylpraecansone B	Muiva, 2012
<i>Tephrosia apollinea</i>	Flavonoid	3,4:8,9-dimethylenedioxypterocarpan (-)-semiglabin  (-)-pseudosemiglabrin (+)-glabratephrin (+)-glabratephrinol appollinine (7-methoxy-8-[3''-(2'',5''-dihydro-5'',5''-dimethyl-2''-oxofuryl)]-flavone	Tarus et al., 2002 Waterman and Khalid, 1980

		lanceolatin-A (+)-apollineanin (-)-semiglabin (-)-semiglabinol	Hisham et al., 2006
<i>Tephrosia barbiger</i>	Flavonoid	isopongaflavone barbigerone	Vilain, 1983 Vilain, 1980
<i>Tephrosia bidwilli</i>	Flavonoid	(-)-6aR; 11aR-maackiain	Ingham and Markham, 1980
		(-)-6aS; 11aS-pisatin (-)-6aR; 11aR-4-methoxy- maackiain tephrocarpin acanthocarpin isopongaflavone	
<i>Tephrosia bracteolata</i>	Flavonoid		Khalid and Waterman, 1981
		trans-tephrostachin trans-anhydrotephrostachin obovatin tephcalostan 7-O-methylglabranin kaempferol 3-O- $\beta$ -D-glucopyranoside (2S)-5-hydroxy-7,4'-di-O-(gamma, gamma- dimethylallyl) flavanone 6-hydroxy-E-3-(2,5-dimethoxy benzylidene)-2',5'-dimethoxyflavanone tephrowatsin C afromosin kaempferol 3-O- $\beta$ -D-glucopyranoside calophione A	Hari Kishore et al., 2003
<i>Tephrosia calophylla</i>	Coumestan Flavonoid		Reddy et al., 2009
		1-(6'-Hydroxy-1',3'-benzodioxol-5'-yl)-2- (6"-hydroxy-2"-isopropenyl-2",3"-dihydro- benzofuran-5"-yl)-ethane-1,2-dione	
	Benzil		Ganapaty et al., 2009b
	Coumestan	tephcalostan B tephcalostan C tephcalostan D	
<i>Tephrosia candida</i>	Flavonoid	candidol candidone ovalichalcone dehydrorotenone candidin pongachin flemichapparin-B $\beta$ -sitsterol caffeic acid 12a-hydroxyrotenone tephrosin amorpolone 6a,12,-dehydodeguelin 12a-hydroxy- $\beta$ -toxicarol deguelin $\alpha$ -toxicarol 6a,12a-dehydrodeguelin 12a-hydroxy- $\alpha$ -toxicarol, 6a 12a-dehydro- $\alpha$ -toxicarol 6a,12a-dehydro- $\beta$ -toxicarol dehydrodihydrorotenone tephrospirolactone tephrospiroketone I tephrospiroketone II demethylapollinin 7-O- $\beta$ -D- glucopyranoside apollinin glabatephrin semiglabin	Dutt and Chibber, 1983 Roy et al., 1986  Parmar et al., 1988  Roy et al., 1987 Parmar et al., 1988  Parmar et al., 1988  Kole et al., 1992 Parmar et al., 1988 Andrei et al., 1997    Roy et al., 1987 Andrei et al., 2002
	Sterol Acid Rotenoid		
<i>Tephrosia cinerea</i>	Flavonoids and Phenolics		Maldini et al., 2011

		pseudosemiglabrin	
		3'-O-methyl- methylquercetin	
		3,7-di-O-rhamnopyranoside	
		kaempferol 3,7-di-O-rhamnopyranoside	
		quercetin 3,7-di-O-rhamnopyranoside	
		3-O- $\beta$ -glucopyranosylquercetin 7-O- $\alpha$ -rhamnopyranoside	
		3-O- $\beta$ -xylopyranosylquercetin 7-O- $\alpha$ -rhamnopyranoside	
		3-O- $\alpha$ -arabinopyranosylquercetin 7-O- $\alpha$ -rhamnopyranoside	
		5-O-methylgenistein 7-O- $\beta$ -D-glucopyranoside	
		quercetin 3-O- $\beta$ -glucopyranoside	
		quercetin 3-O- $\alpha$ -rhamnopyranoside	
		kaempferol	
		7-O-methylquercetin	
		cineroside A	
	Sesquiterpene	caryophyllene oxide	Arriaga et al., 2008
		teclenone B	
		(1 $\beta$ ,7R*)-opposit-4(15)-ene-1,7-diol	
	Lignan	pinoresinol	
<i>Tephrosia crassifolia</i>	Flavonoid	crassifolin	Gómez-Garibay et al., 1999
<i>Tephrosia egreria</i>	Terpenoid	crassichalcone	
		geijerene	Arriaga et al., 2005
		pregeijerene	
<i>Tephrosia elata</i>	Rotenoid	Dehydrorotenone.	Arriaga et al., 2009b
	Flavonoid	isopongaflavone	Bentley et al., 1987
		tephrosin	
		8-(3,3-dimethylallyl)-5,7- dimethoxy flavanone	Lwande et al., 1985a
		obovatin methyl ether	
		warangalone	
		elatadihydrochalcone	Muiva, 2012
		obovatachalcone	
		(S)-elatadihydrochalcone	Muiva et al., 2009
		obovatachalcone	
		obovatin	Muiva et al., 2009; Muiva, 2012
		obovatin methyl ether	Muiva et al., 2009; Muiva, 2012
	Pterocarpan	(+)-pisatin	Lwande et al., 1985a
		(-)- maackiain	
	Rotenoid	deguelin	Muiva et al., 2009; Muiva, 2012
<i>Tephrosia elongata</i>	Flavonoid	rotenone	Muiva, 2012
<i>Tephrosia emoroides</i>	Flavonoid	elongatin	Smalberger et al., 1975
		emoroidenone	Machocho et al., 1995
		emoroidone	
		emoroidocarpin	
		5-methoxyisolonchocarpin	
<i>Tephrosia falciformis</i>	Flavene	hildegardtene	
	Flavonoid	falciformin	Khan et al., 1986
		7-hydroxy-8-( $\gamma$ , $\gamma$ -dimethylallyl)flavanone	
<i>Tephrosia fulvinervis</i>	Alcohol	triacontanol	Khan et al., 1984
	Flavonoid	fulvinervin C	Venkataratnam et al., 1986
		fulvinervin A	Venkataratnam et al., 1986; Venkata et al., 1985b
		fulvinervin B	Venkata et al., 1985b
	Rotenoid	$\alpha$ -toxicarol	Dagne et al., 1989

		deguelin munduserone cis-12a-hydroxymunduserone (-)-maackiain	
<i>Tephrosia hamiltonii</i>	Pterocarpan Flavonoid	5, 7-dimethoxy-8-(2, 3-epoxy-3-methylbutyl)-flavanone pongamol flemichapparin-B flemichapparin-C	Falak and Shoeb 1987 Rajani and Sarma, 1988
<i>Tephrosia hildebrandtii</i>	Coumestone Pterocarpan	2-methoxy-3,9-dihydroxy coumestone hildecarpidin	Lwande et al., 1987
	Flavonoid	hildecarpin trans-tephrostachin	Lwande et al., 1985b Lwande et al., 1986
<i>Tephrosia hookeriana</i>	Flavonoid	trans-anhydrotephrostachin hookerianin	Prabhakar et al., 1996; Vanangamudi et al., 1997b
		(-)-semiglabin lanceolatin A. tephrorianin	Vanangamudi et al., 1997b
<i>Tephrosia lanceolata</i>	Flavonoid	rutin rutin	Rangaswami and Rao, 1955
<i>Tephrosia leiocarpa</i>	Flavonoid	tephroleocarpin A tephroleocarpin B	Quijano and Rios, 1991
<i>Tephrosia lupinifolia</i>	Flavonoid	lupinifolin lupinifolinol	Smalberger et al., 1974
<i>Tephrosia madrensis</i>	Flavonoid	5,7-dimethoxy-8-prenylflavan	Gómez et al., 1983
<i>Tephrosia major</i>	Flavonoid	2',6'-dihydroxy-3'-prenyl-4'-methoxy- $\beta$ -hydroxychalcone quercetin	Gomez-Garibay et al., 2002
	Sterol	$\beta$ -sitosterol stigmasterol	
<i>Tephrosia maxima</i>	Triterpene Flavonoid	lupeol maxima isoflavone A maxima isoflavone A maxima isoflavone B	Venkata et al., 1994 Rao et al., 1984a Venkata and Sree Rama, 1985a Rao et al., 1984a
		maxima isoflavone C maxima isoflavone D maxima isoflavone E maxima isoflavone F maxima isoflavone G maxima isoflavone H	
		maxima isoflavone J	Venkata and Sree Rama, 1985a
<i>Tephrosia multijuga</i>	Flavonoid	maxima isoflavone T multijuginol multijugin	Murthy and Rao, 1985; Venkata et al., 1994 Venkata et al., 1994 Vleggaar et al., 1975
<i>Tephrosia nubica</i>	Flavonoidal glycoside	kaemferol 3,7-dirhamnoside	Sharaby and Ammar, 1997
	Flavonoid	quercetin 3-galactoside 7-rhamnoside quercetin 3,7-dirhamnoside semiglabin pseudosemiglabin apollinine lanceolatin A	
	Rotenoid	rotenones	
<i>Tephrosia</i>	Rotenoid	deguelin dihydrostemonal	Dagne et al., 1989

<i>pentaphylla</i>		9-demethyldihydrostemonal 6-acetoxydihydrostemonal villosin sumatrol rotenone cis-12a-hydroxyrotenone 6-hydroxyrotenone $\alpha$ -toxicarol	
<i>Tephrosia polyphylla</i>	Flavonoid Flavonoid	obovatin 4'-demethyltoxicarol isoflavone toxicarol isoflavone 7-methylglabranin	Dagne et al., 1992
<i>Tephrosia procumbens</i>	Rotenoid	rotenone	Venkataratnam et al., 1987
	$\beta$ -diketone	sumatrol praecansone A praecansone B	
	Flavonoid	obovatin 7-ethoxy-3,3',4'-trihydroxyflavone; fisetin 7-ethyl ether	
<i>Tephrosia pumila</i>	Flavonoid	7,4'-dihydroxy-3'-methoxyisoflavone pumilaisoflavones A pumilaisoflavones B pumilaisoflavones C pumilaisoflavones D pumilanol tephrinone $\beta$ -hydroxychalcone Praecansone-A.	Yenesew et al., 1989 Ganapaty et al., 2008b Dagne et al., 1988
	Rotenoid	rotenone	Ganapaty et al., 2008b
	Triterpene	lupeol	
<i>Tephrosia purpurea</i>	Sterol Flavonoid	stigmasterol tephrosin pongaglabol semiglabin purpuritenin purpureamethide pongamol karanjin lanceolatin B	Ahmad et al., 1999 Sinha et al., 1982
		(+)-tephrorins A (+)-tephrorins B (+)-tephrosone purpurenone (+)-purpurin	Sinha et al., 1982; Chang et al., 1997 Chang et al., 2000 Rao and Raju, 1984b Rao and Raju, 1984b; Chang et al., 1997
		(-)-purpurin dehydroisoderricin (-)-maackiain pseudosemiglabrin (-)-semiglabin terpurinflavone pongamol	Juma et al., 2011 Parmar et al., 1989; Chang et al., 1997
		(-)-isolonchocarpin 7,4'-dihydroxy-3',5'-dimethoxyisoflavone (+)-tephropurpurin (-)-3-hydroxy-4-methoxy-8,9-methylenedioxypterocarpan (-)-medicarpin 3'-methoxydaidzein	Rao and Raju, 1979 Chang et al., 1997

		desmoxyphyllin B 3,9-dihydroxy-8-methoxycoumestan isoglabratephrin tephropurpulin A quercitin rutin	Hegazy et al., 2009 Jain et al., 2009
	Ester	stigmast-5, 22-dien-34, 21diol-34, 21-dihexadecanoate	Sharma et al., 2008
	Neoflavonoid glycoside Sterol	serratin 7-O-[ $\beta$ -D-glucopyranosyl-(1 $\rightarrow$ 4)-O- $\beta$ -D-galactopyranoside $\beta$ -sitosterol	Saxena and Choubey, 1997 Chang et al., 1997; Parmar et al., 1989
<i>Tephrosia quercetorum</i>	Acid Flavonoid	spinasterol- $\alpha$ ursolic acid quercetols A	Gómez-Garibay et al., 1988
<i>Tephrosia semiglabra</i>	Flavonoid	quercetols B quercetols C glabratephrin semiglabrinol semiglabrin	Vleggaar et al., 1978 Smalberger et al., 1973
<i>Tephrosia sinapou</i>	Flavonoid	toxicarine 7-O-methylglabranine tephrowatsin A quercetol B flamichapparin B	Martinez et al., 2012
	Coumarin Rotenoid	2,3-dihydro-p-coumaric acid tephrosin rotenolone deguelin 6-oxo-6a,12a-dehydrodeguelin 6-oxo-6a,12a-dehydro- $\alpha$ -toxicarol 6a,12a-dehydrorotenone rotenonone villosone	
<i>Tephrosia spinosa</i>	Flavonoid	spinochalcone C spinoflavanones A spinoflavanones B fulvinervin A 3',5'-diisopentenyl-2',4'-dihydroxychalcone tephrospinosin spinochalcones A spinochalcones B flemistrictin A	Rao and Prasad, 1992 Sharma and Rao, 1992 Rao and Prasad, 1992
<i>Tephrosia tepicana</i>	Flavonol glycoside	eupalitin 3-O-b-D-galactopyranoside	Vanangamudi et al., 1997a; Chakradhar et al., 2005
<i>Tephrosia tinctoria</i>	Flavonoid	tepicanol A	Gómez-Garibay et al., 1997
	Flavonoid	5,7-di-O-prenylbiochanin A 7-O-methylglabranin tephrowatsin C flemichapparin B 2-hydroxy tephrosin tephrinone lupinifolin 7-O-methyl glabranin	Khalivulla et al., 2008 Ganapaty et al., 2009 Ganapaty et al., 2010
	Rotenoid	rotenone dehydrodeguelin	
	Sterol	stigmasterol	
<i>Tephrosia toxicaria</i>	Acid Flavonoid	betulinic acid iso-obovatin obovatin	Vasconcelos et al., 2009 Vasconcelos et al., 2009;

		6a,12a-dehydro- $\beta$ -toxicarol	Jang et al., 2003
		6a,12a-dehydro- $\alpha$ -toxicarol	
		$\alpha$ -toxicarol	
		toxicarol	Clark, 1930
		(2S)-5-hydroxy-7-methoxy-8-[(E)-3-oxo-1-	Jang et al., 2003
		butenyl]flavanone	
		isoliquiritigenin	
		genistein	
		chrysoeriol	
		sumatrol	
	Rotenoid	4',5'-dihydro-11,5'-dihydroxy-4'-	
		methoxytephrosin	
		11-hydroxytephrosin	
	Coumarin	marmesin	
	Triterpene	lupenone	
	Ester	benzyl benzoate	
		benzyl trans-cinnamate	
<i>Tephrosia tunicata</i>	Flavonoid	tunicatachalcone	Andrei et al., 2000
<i>Tephrosia uniflora</i>	Flavonoid	elongatin	Abreu and Luis, 1996
	Rotenoid	12a-hydroxyrotenone	
	Sterol	$\beta$ -sitosterol	
		stigmasterol	
<i>Tephrosia viciodes</i>	Flavonoid	enantiomultijugin	Gómez-Garibay et al., 1992
<i>Tephrosia villosa</i>	Flavonoid	(2S)-5,4'-dihydroxy-7-O-[(E)-3,7-dimethyl-	Madhusudhana et al., 2010
		2,6-octadienyl]flavanone	
		(2S)-5,4'-dihydroxy-7-O-[(E)-3,7-dimethyl-	
		2,6-octa-dienyl]-8-C-[(E)-3,7-dimethyl-2,6-	
		octadienyl]flavanone	
		7-O-methylglabranin	
		tephcalostan	
		12a-dehydro-6-hydroxysumatrol	
		7-methylglabranin	Jayaraman et al., 1980
		villosin	David Krupadanam et al., 1997
		villosone	
		villol	
		villinol	
		tephrinone	Rao and Srimanarayana, 1981
	Triterpenoid	lupenone	Prashant and Krupadanam 1993
	Triterpene	lupeol	Ganapaty et al., 2008a
	Sterol	stigmasterol	
	Rotenoid	12a-dehydro-6-hydroxysumatrol	Prashant and Krupadanam 1993
		rotenone	Ganapaty et al., 2008a
		dehydrorotenone	
		6a,12a-dehydro,2,3,6- trimethoxy-8-(3',3'-	Prashant and Krupadanam, 1993
		dimethylallyl)-9,11dihydroxy rotenone	
		12a-hydroxy toxicarol	
<i>Tephrosia viridiflora</i>	Flavonoids	viridiflorin	Gómez et al., 1985
<i>Tephrosia vogelii</i>	Sesquiterpene	(1 $\beta$ ,6 $\alpha$ ,10 $\alpha$ )-guai-4(15)-ene-6,7,10-triol	Wei et al., 2009
	Lignan	(+)-lariciresinol 9'-stearate	
	Rotenoid	deguelin	Kalume et al., 2012; Delfel et al., 1970; Gills, 1992
		tephrosin	Gills, 1992
		toxicanol	
		tephrosal	
	Flavonoid	quercetin	
<i>Tephrosia</i>	Flavonoid	tephrowatsin A	Gómez et al., 1985




watsoniana

*Tephrosia woodii*


Flavonoid

tephrowatsin B  
tephrowatsin C  
tephrowatsin D  
tephrowatsin E  
oaxacacin  
mixtecacin


Dominguez et al., 1983




(-)-6aR,11aR,4-Methoxy Maackiain




(-)-6aR,11aR-Maackiain




(-)-Acanthocarpan




(-)-Medicarpin




(1B,7R)-opposit-4(15)-ene-1,7-diol




3-Methoxy Daidzein




5-Methoxy Isolonchocarpin




7,4-dihydroxy 3-methoxy isoflavone




7-O-Methyl Quercetin




12a-hydroxyrotenone




3,4,8,9-dimethylene dioxy pterocarpan




Afroformosin




Amorfolone




Benzyl Benzoate




Falciformin




Benzyl Cinnamate




Calophione A




Calophyllene Oxide




Candidin




Candidol




Candidone




Chrysoeriol




Dehydrotetrahydro Rotenone




Dihydro P-Coumaric Acid




Dehydro Rotenone




Demethyl Apollinin 7-O-Beta-D Glucopyranoside




Geijerene




Demethyl Praecansone B




Desmoxyphyllin B




Dihydrostemonal











Elatadihydrochalcone




Elongatin




Emoroidenone







Genistein




Hildecarpidin




Hildecarpin




Hookerianin




Isoglabratephrin




Isoliquiritigenin




Isopongaflavone




Kaempferol 3,7 dirhamnoside




Kaempferol




Karanjin




Lanceolatin A




Lupinifolin




Lupinifolinol




Marmesin




Maxima isoflavone C




Maxima isoflavone D




Maxima isoflavone G




Multijugin




Multijuginol




Munduserone




2',6'-dihydroxy-3'-prenyl-4'-methoxy-β-hydroxychalcone




Oaxacacin




Obovatachalcone




Obovatin methyl ether




Ovalichalcone




Pinoresinol




Pisatin




Pongachin




Pongaglabol




Pongamol




Praecansone B




Pumilaisoflavone A




Pumilaisoflavone B




Pumilaisoflavone C




Pumilaisoflavone D




Pumilanol




Purpureamethide



Purpuritenin




Quercetin 3 galactoside 7 rhamnoside




Quercetin 3,7 dirhamnoside







Quercetol A




Quercetol B




Quercetol C




Rotenolone




Semiglabinol




Spinasterol




Spinochalcone A




Spinochalcone B




Spinochalcone C




Spinoflavanone A




Stigmasterol




Sumatrol




Teclenone A




Teclenone B




Tephcalostan B




Tephcalostan C




Tephcalostan D




Tephcalostan A




Tephrocarpin




Tephroleocarpin A




Tephroleocarpin B




Tephrorianin




Tephrospiroketone 1




Tephrospiroketone 2




Tephrospirolactone




Tephrowatsin A




Tephrowatsin B




Tephrowatsin C




Tephrowatsin D




Tephrowatsin E




Trans tephrostachin




Trans anhydrotrophostachin




Tunicatachalcone




Ursolic Acid




Villinol




Villosone




Viridiflorin




Villol




Warangalone




5,7 dimethoxy 8 prenylflavan




(-)- Pseudoemiglabrin




(+)- Glabratephrinol




7-O- Methyl Glabranin




(+)- Glabratephrin




Betulinic Acid




Apolinine




Barbigerone




Caffeic acid




Cinoside A




Deguelin




Flemichapparin A




Fulvinervin A




Hildecarpin




Isopongaflavone




Lupeol




Maxima isoflavone A




Millettone




Obovatin




Praecansone A




Purpurin




Quercetin




Rotenone




Tephrosin




Rutin



Semiglabin



b-sitosterol



Some of the compounds isolated have been studied for their pharmacological actions (see table 2). The activities reported include anticancer, antiplasmodial, larvicidal, and others. There are many compounds in Table 1 whose activities are not studied under the genus *Tephrosia* but if we look into the literature, we find their presence in other genera and their activities determined for example Pseudosemiglabrin, Flemichapparin, Caryophyllene oxide, deguelin, pongamol, lupeol, possess platelet aggregation antagonism, antifungal, anticancer, anticonvulsant, and antiinflammatory respectively (Pirrung and Lee, 1995; Gahlot et al., 2012; Yang et al., 2000; Udeani et al., 1997; Basu et al., 1994; Geetha and Varalakshmi, 2001). Caffeic acid and rutin, which are also found in *Phyllanthus sellowianus* have analgesic activity (Calixto et al., 1998).

Table 2. Pharmacological actions of isolated compounds from genus *Tephrosia*.

Species	Compound	Activity	Reference
<i>Tephrosia calophylla</i> <i>Tephrosia candida</i>	calophione A	Cytotoxic	Ganapaty et al., 2009
	candidone	Cytotoxic	Ganapaty et al., 2009; Roy et al., 1986
	pongachin		Ganapaty et al., 2009; Parmar et al., 1988
<i>Tephrosia elata</i>	candidachalcone	Estrogenic activity	Hegazy et al., 2011
	tephrosin	Antifeedant	Bentley et al., 1987
	isopongaflavone rotenone	Larvicidal, antifeedant	Muiva, 2012; Bentley et al., 1987
<i>Tephrosia emoroides</i> <i>Tephrosia ergeria</i> <i>Tephrosia hildebrandtii</i>	(S)-elatadihydrochalcone	Antiplasmodial	Muiva, 2012
	obovatin methyl ether		Muiva, 2012
	praecansone	Antifeedant	Machocho et al., 1995
	emoroidenone	Antioxidant, larvicidal	Arriaga et al., 2009a
<i>Tephrosia hildebrandtii</i>	dehydrorotenone	Insect antifeedant	Lwande et al., 1985
	hildecarpin	Antifungal	
<i>Tephrosia pulcherrima</i> <i>Tephrosia pumila</i> <i>Tephrosia purpurea</i>	pulcherrimin	Cytotoxic	Ganapaty et al., 2009
	pumilanol	Antiprotozoal	Ganapaty et al., 2008
	(+)-tephrorin A	Cancer chemopreventive activity	Chang et al., 2000



	(+)-tephrosin B		
	(+)-tephrosone		
	7,4'-dihydroxy-3',5'- dimethoxyisoflavone		Chang et al., 2000
	(+)-tephropurpurin		
	(+)-purpurin		
	pongamol		
	lanceolatin B		
	(-)-maackiain		
	(-)-3-hydroxy-4-methoxy - 8, 9- methylenedioxy		
	pterocarpan		
	(-)-medicarpin		
	terpurinflavone	Antiplasmodial	Juma et al., 2011
<i>Tephrosia spinosa</i>	eupalitin-3-O- $\beta$ -D-glucoside	Anti-inflammatory	Chakradhar et al., 2005
<i>Tephrosia tinctoria</i>	2-hydroxy tephrosin	Antiplasmodial	Ganapaty et al., 2009
	tephrinone		
<i>Tephrosia toxicaria</i>	(2S)-5-hydroxy-7-methoxy- 8-[(E)-3-oxo-1- butenyl]flavanone	Cancer chemopreventive activity	Jang et al., 2003
	4',5'-dihydro-11,5'- dihydroxy-4'- methoxytephrosin		
	isoliquiritigenin		
	genistein		
	chrysoeriol		
	obovatin	Antioxidant	Vasconcelos et al., 2009
	toxicarol	Fish poison	Clark, 1930
	$\alpha$ -toxicarol	Larvicidal	Vasconcelos et al., 2009
<i>Tephrosia vogelii</i>	deguelin	Larvicidal	Muiva, 2012; Kalume et al., 2012

It can be concluded from the above discussion that if, for example, any species of genus *Tephrosia* having rutin or caffeic acid as its major component, is not reported for analgesic activity, may possess it. So in this way this methodology can give us a hint or base, which study to be carried on plant.

### Pharmacological profile of plants from genus *Tephrosia*

Several plants of the genus have studied for their medicinal and therapeutic potential. A brief description of the work done so far is given below:

#### *Antioxidant activity*

Only a few species of Genus *Tephrosia* have been studied for their antioxidant activity. In 2007, G.P. Choudhary studied the ethanolic extract of *Tephrosia purpurea* for its antioxidant activity (Choudhary, 2007). The aqueous extract of the whole plant of *Tephrosia purpurea* also showed free radical scavenging activity in DPPH test (Gunjegaonkar et al., 2010). From *Tephrosia egregia* the ethyl acetate and methanol extracts showed high antioxidant activities (Arriaga et al., 2009a). Obovatin, a flavonoid present in *Tephrosia toxicaria* showed significant antioxidant activity of IC<sub>50</sub> 3.370  $\mu$ g/mL. It was also seen that the methanol fraction of the ethanol extract from roots had the highest antioxidant activity (Vasconcelos et al., 2009). *Tephrosia villosa* also possess antioxidant activity due to the presence of 20(29)-lupen-3-one, a compound also identified in *Daedaleopsis tricolor* where it inhibited lipid peroxidation by 6.4% (Prashant and Krupadanam 1993; Kim et al., 2001). The ethanol

ether extract of *Tephrosia vogelii* seeds also showed antioxidant and free radical scavenging activity (Li et al., 2010).

### **Antibacterial activity**

The species from Genus *Tephrosia* have also been studied for their antibacterial activity. *Tephrosia vogelii* was found to possess antimicrobial activity (Wanga et al., 2007). The dichloromethane extract from the roots and leaves was tested against *S. aureus*, *E. coli* and *F. phoseolida*. Hu et al., in 2011 also studied the antimicrobial and bacteriostatic activity of ethanol and aqueous extract from *Tephrosia vogelii* seeds on *E. coli*, *S. aureus* and *S. paratyphi* B. and proved the antibacterial efficacy of the plant to be significant at high doses (Hu et al., 2011). The root extract of *Tephrosia villosa* showed moderate antibacterial and anti fungal activity (Ganapaty et al., 2008a). In another study on *Tephrosia villosa* the fruit, leaf, and root extract showed activity against *C. neoformans*, *E. coli* and *B. anthracis* respectively. The ethanolic twig extract was most active against *C. neoformans* and *S. typhi* (Nondo et al., 2011). In case of *Tephrosia purpurea*, studies have been made on the antimicrobial activity of methanolic extract of *Tephrosia purpurea* roots on *B. subtilis*, *S. aureus*, *M. luteus*, the gram positive bacteria and the gram negative including *E. coli*, *P. aeruginosa*, and *S. typhimurium* (Gupta et al., 2008). In another study on *Tephrosia purpurea*, the roots showed antimicrobial activity against *P. aeruginosa* and no activity against *S. aureus* and *E. coli* (BNLD Rangama et al., 2009). Chinniah et al., in 2009 and Annalakshmi et al., in 2009 proved *Tephrosia purpurea* to have marked activity against *H. pylori*, an agent responsible for GIT ulcers (Chinniah et al., 2009; Annalakshmi et al., 2009). The methanolic leaf extract from *Tephrosia tinctoria* showed activity against *B. subtilis*, *S. marcescens*, and low activity for *B. cereus* and *P. aeruginosa* (Ganapaty et al., 2010). *Tephrosia deflexa* and its isolated compounds were studied for antibacterial activity (Kare et al., 2006). The antibacterial activity of *Tephrosia linearis* has also been reported (Ratsimamanga et al., 1994).

### **Antifungal activity**

In our literature survey, we found less work on antifungal activity of species from Genus *Tephrosia*. Only three species are known to possess antifungal potential. The methanolic extract of *Tephrosia purpurea* showed significant activity against *A. niger* and *C. albicans* (Gupta et al., 2008). *Tephrosia hildebrandtii* showed antifungal activity against *C. cucumerinum*. The activity was found to be related to a chemical compound isolated from its roots (see table 2) (Lwande et al., 1985b). *Tephrosia tinctoria* also showed activity against *A. niger*, *C. albicans*. The methanolic extract was found to be more active against the aforementioned organisms. However the methanolic extract showed no activity against *S. cerevisiae* (Ganapaty et al., 2010).

### **Antiprotozoal and anti plasmodial activity**

Extract from the seed pods of *Tephrosia elata* showed antiplasmodial activity (Muiva et al., 2009; Muiva, 2012). Flavonoid extracted from the roots of *Tephrosia pumila* also showed activity against *L. donovani*, *T. b. rhodesiense* and *T. cruzi* (Ganapaty et al., 2008b). Isolated flavonoids from the root of *Tephrosia tinctoria* were studied for antiprotozoal and

antiplasmodial activities against *T. b. rhodesiense*, *T. cruzi*, *L. donovani*, and *P. falciparum* (Ganapaty et al., 2009a). Ganapaty, also studied the antiprotozoal activity of three *Tephrosia* species, namely, *T. pulcherrima*, *T. pumila*, and *T. calophylla* on *Leishmania*, *Trypanosoma* and *Plasmodium* parasites (Ganapaty et al., 2009c). Chloroquine sensitive and chloroquine resistant strains of *P. falciparum* were inhibited by the extracts from the stem of *Tephrosia purpurea* with IC<sub>50</sub> values of 10.47 ± 2.22 µg/ml and 12.06 ± 2.54 µg/ml, respectively (Juma et al., 2011). *Tephrosia purpurea* has also been studied for antileishmanial activity in hamsters and Indian langular monkeys infected by *L. donovani* (Sharma et al., 2003).

### **Anti pyretic and Anti inflammatory activity**

In 2010, Sandhya et al., studied the anti inflammatory activity of two species of *Tephrosia* namely *Tephrosia maxima* and *Tephrosia purpurea* by HRBC membrane stabilizing method. Both plants showed almost equal activity at doses of 500ug/ml. *Tephrosia maxima* giving 79.49% and *Tephrosia purpurea* giving 79.01% protection (Sandhya et al., 2010). Another study on *Tephrosia purpurea* root extracts showed its antipyretic and anti inflammatory activity (Valli et al., 2011). The methanolic extract of *Tephrosia vogelii* showed significant analgesic and anti-inflammatory activity in mice and rats using hot plate method and egg albumin induced oedema respectively (Audaud et al., 2009). The root extract of *Tephrosia sinapou* showed to possess significant anti-inflammatory activity. The extract reduced inflammatory leukocyte recruitment, oxidative stress and other parameters involved directly or indirectly to the process of inflammation (Martinez et al., 2012). *Tephrosia spinosa* also showed anti inflammatory activity in an experimental model of carrageenin induced paw edema. The standard drug used was indomethacin (Chakradhar et al., 2005). The antipyretic activity of *Tephrosia bracteolata* has also been reported (Onaolapo et al., 2009).

### **Anticancer and cytotoxic activity**

Cytotoxicity of some chemical compounds found in *Tephrosia calophylla* and *Tephrosia candida* have been studied using different cell lines (Ganapaty et al., 2009a; Ganapaty et al., 2009b; Roy et al., 1986; Parmar et al., 1988). The cytotoxicity of *Tephrosia pulcherrima* and *Tephrosia pumila* has also been studied by Ganapaty et al., in 2009 using HT-29 and RAW cell lines (Ganapaty et al., 2009c). In 2011 Kishore et al., mentioned *Tephrosia purpurea* containing an important chemical, B-sitosteol having anticancer and cancer protective activities against prostatic, breast and colonic carcinomas. In addition to the aforementioned activities of B-sitosterol, it is also an antioxidant and has significant effect on hypercholesterolemia and BPH (Kishore and Roy, 2011). In another study the anticarcinogenic activity of *Tephrosia purpurea* extract was tested in an experimental model of hepatocarcinoma in rats. The extract showed significant cancer chemoprevention (Hussain et al., 2012). Shanmugapriya et al., also studied the anticarcinogenic potential of *Tephrosia purpurea* in HELA cervical cancerous cell line. Different extracts were tested out of which ethyl acetate produced the most potent effect (Shanmugapriya et al., 2011). In a study by Subhadra, three species namely, *Tephrosia calophylla*, *Tephrosia maxima* and *Tephrosia purpurea* showed significant cytotoxic activity out of which *Tephrosia calophylla* showed the maximum activity (Subhadra et al., 2011). The ethanolic fruit and root extract of *Tephrosia villosa* showed toxicity to brine shrimp whereas the extract from leaves and twigs was found to be non toxic (Nondo et al., 2011). The ethyl acetate extract from stems of *Tephrosia toxicaria* possess

flavonoids having cancer chemopreventive activities (Jang et al., 2003). The flavonoids extracted from *Tephrosia tinctoria* possess cytotoxic activity tested in Cell line L-6 (Rat skeletal muscle myoblasts) (Ganapaty et al., 2009a). *Tephrosia calophylla* was also found to possess anticancer activity. The root extract inhibited growth and induced apoptosis in the human breast carcinoma (Adinarayana et al., 2009). *Tephrosia vogelii* root and leaf extract was found to be toxic to brine shrimps at doses of LC50: 0.960; 0.958 µg/ml, respectively (Wanga et al., 2007).

### **Hepatoprotective activity**

In 2011, Shah et al., studied the hepatoprotective effect of *Tephrosia purpurea* in CCl<sub>4</sub> induced hepatotoxicity in rats. The ethyl acetate fraction at doses of 50mg/kg was found to be effective and comparable to silymarin (Rajal Shah et al., 2011). The hepatoprotective effect of *Tephrosia calophylla* has also been reported (Adinarayana et al., 2011).

### **Animal feed**

In an effort to find new and cheap sources of food for animals, several species of genus *Tephrosia* have been studied. The nutritive value of three species of *Tephrosia*, namely, *Tephrosia candida*, *Tephrosia bracteolata*, and *Tephrosia linearis* have been studied (Babayemi et al., 2003). According to Babayemi and Bamikole, a mixture of *Tephrosia candida* leaves and guinea grass can serve as a good animal feed. The mixture has an additional benefit of low methane production upon fermentation (Babayemi and Bamikole. 2006b). *Tephrosia bracteolata* can serve as a good diet in laying hens both from nutritive and economic aspect (Akande et al., 2008). *Tephrosia vogelii*, *Tephrosia candida*, and *Tephrosia purpurea* can also be a good addition in the diet of ruminants (Mbomi et al., 2011). The use of *Tephrosia candida* and *Tephrosia bracteolata* in goats has also been established (Babayemi and Bamikole 2006a). A study on *Tephrosia candida* seeds has also been reported (Babayemi and Bamikole 2007).

### **Larvicidal, insecticidal and antifeedant activity**

Different species from the genus have been studied for larvicidal, insecticidal, and antifeedant activities. There is an extensive work on the study of *Tephrosia* as an agent to control the population of insects harmful to animals and plants.

The hexane extract from *Tephrosia egregia* showed potent larvicidal activity against aedes aegypti (Arriaga et al., 2009a). The whole plant extract of *Tephrosia purpurea* was tested for its larvicidal activity against the larvae of *Culex quinquefasciatus*. The extract showed 100% mortality in very small doses suggesting its beneficial use in controlling the mosquito reproduction (Deepak Kumar et al., 2012). The extracts of *Tephrosia vogelii* also possess larvicidal activity and therefore can be used to control mosquitoes (Matovu and Olila. 2007). The ethanolic extract of roots, leaves, fruit and twigs of *Tephrosia villosa* showed significant activity against *C. quinquefasciatus* larvae (Nondo et al., 2011). The ethanol extract from roots, stems, leaves, and pods and some fractions of *Tephrosia toxicaria* were tested for larvicidal activity with the larvae of *Aedes aegypti*. The ethanolic root extract, hexane and chlor-

oform fractions had (LC<sub>50</sub> 47.86 ppm) (LC<sub>50</sub> 23.99 ppm) and (LC<sub>50</sub> 13.80 ppm) respectively (Vasconcelos et al., 2009). *Tephrosia nyikensis* have been reported to possess larvicidal activity on Anopheles mosquito's larvae (Wanjala et al., 2006). The oil obtained from *Tephrosia cinerea* showed larvicidal activity against *Aedes aegypti* larvae (Arriaga et al., 2008). The chloroform and methanol extracts of *Tephrosia nubica* were tested against *Spodoptera littoralis* and *Agrotis ipsilon*. The population of the pests was reduced due to the effect of the extract on all the stages of growth (Sharaby and Ammar, 1997). *Tephrosia vogelii* leaf extract was found to be effective in controlling ticks, an important insect and ectoparasite (Gadzirayi et al., 2010). *Tephrosia magropoda* is also reported to have insecticidal properties (Tatteksfield and Gimingham, 1932). In 2012, Kalume et al., reported the acaricidal activity of leaf extracts of *Tephrosia vogelii* on tick *Rhipicephalus appendiculatus* and mentioned its advantage of being economical than synthetic compounds (Kalume et al., 2012). The insecticidal property of *Tephrosia purpurea* whole plant was tested against *Callosobruchus maculatus*, the pest on *Phaseolus mungo* (Diwan and Saxena, 2010). In 1992, Kole et al., isolated a rotenoid, amorpholone from *Tephrosia candida* having potent insecticidal properties (Kole et al., 1992). *Tephrosia elata* showed significant antifeedant activity against *M. testulalis*, *S. exempta* and *E. sacchariana*. The antifeedant activity is attributed to the presence of rotenoid compounds (Bentley et al., 1987). The larvicidal activity from seed pods of *Tephrosia elata* and *Tephrosia aequilata* has also been studied by Muiva, against the larvae of *Aedes aegypti* (Muiva, 2012). Antifeedant activity of flavonoids from *Tephrosia emoroides* was tested against *Chilo partellus*, a very destructive pest of maize. Emoroidenone, a flavonoid isolated showed strong feeding deterrence of 66.1% against the larvae at a dose of 100 µg (Machocho et al., 1995). The roots of *Tephrosia hidebrandtii* also possess antifeedant activity against the pest, *Maruca testulalis* (Lwande et al., 1985).

### **Antidiabetic activity**

The aqueous seed extract of *Tephrosia purpurea* showed significant antihyperglycemic activity in streptozotocin induced diabetic rats (Pavana et al., 2009). The ethanolic extract of from *Tephrosia villosa* leaves showed reduction in glucose level and pancreatic cell regeneration in alloxan induced diabetes in rats (Ahmad et al., 2009). Balakrishnan et al., also reported antidiabetic activity of extract from root of *Tephrosia villosa* (Balakrishnan et al., 2007).

### **GIT activity**

*Tephrosia vogelii* leaf extract exerted a stimulant effect on the GIT smooth muscles. This was demonstrated by the contraction of the ileum isolated from guinea pig hence showing the purgative property of the plant (Dzenda et al., 2008b). Aqueous extract of *Tephrosia purpurea* root showed gastric ulcer healing and cytoprotective activities (Deshpande and Shah 2008). The extract of *Tephrosia calophylla* leaves showed significant antiulcer and cytoprotective activity at doses of 50mg/kg and 100mg/kg (Divya, et al., 2011).

### **Antihyperlipidemic effect**

The antihyperlipidemic effect of *Tephrosia calophylla* has been studied in wistar albino rats (Mohan, 2011). The leaf extract of *Tephrosia purpurea* showed antihyperlipidemic

activity in an experimental model of diabetic rats (Pavana et al., 2007). Akhtar et al., also studied *Tephrosia purpurea* for the same purpose and found a significant reduction in all the parameters (Akhtar et al., 2011).

### **Toxicity**

*Tephrosia purpurea* extract was evaluated by Talib et al., in 2012 for its toxicity in rodents. A dose up to 2000mg/kg was well tolerated in the acute toxicity studies whereas in sub acute toxicity studies, a dose 200mg/kg and 400 mg/kg showed no significant change in any of the parameters thus concluding that the plant is safe for use in treatment of different diseases (Talib Hussain et al., 2012). *Tephrosia toxicaria* used as a fish poison was studied by Clark in 1930. A compound, Toxicarol was identified as the major component (Clark, 1930). The toxicity of *Tephrosia vogelii* was reported on mice. The signs were similar to those associated with the toxicity from rotenone. The LD<sub>50</sub> of leaf extract calculated was 134.16 mg/kg (Dzenda et al., 2008a). The chloroform extract of *Tephrosia tinctoria* leaves exhibited significant piscicidal activity compared to methanolic extract in gold fish (Ganapaty et al., 2010). Toxic hepatopathy was reported in sheep grazing on *Tephrosia cinerea*. The disease was also experimentally induced in the sheep in order to confirm the results (Santos et al., 2007). *Tephrosia apollinea* was also found to be toxic in a study on goats (Suliman et al., 1982). The toxicity of *Tephrosia bracteolata* has also been studied (Onaolapo et al., 2009). In a study on mice Cai et al., found *Tephrosia candida* to be safe and no significant signs of toxicity were observed (Cai et al., 2010).

### **Miscellaneous activities**

The root extract of *Tephrosia purpurea* showed xanthine oxidase inhibitory activity compare with standard, Allopurinol (Nile and Khobragade, 2011). Patel et al., studied the effect of *Tephrosia purpuria* on polycystic ovary syndrome (PCOS) in rats. (PCOS) was induced by the administration of Letrozole. The dried seed powder given orally showed normalization in the estrous cycle and reduction in the weight of the reproductive system as well as of the ovary (Patel and Thakor, 2012). Kumar et al., found *Tephrosia purpuria* to be effective anxiolytic agent and comparable to the standard drug, Diazepam. The hydroalcoholic extract at a dose of 200mg/kg and 400mg/kg orally was administered to mice in different maze models in the study (Kumar, et al., 2011). The acetylcholinesterase inhibitory activity of *Tephrosia purpurea* and neurobehavioral studies were made on zebra fish, a model for the study of neurodegenerative activities (Kannan and Vincent, 2012). *Tephrosia purpurea* has also been proved for its antiepileptic effect (Asuntha et al., 2010). Lodhi et al., studied the flavonoidal extract of *Tephrosia purpurea* and proved its potential for healing burn wounds. This activity is supposed to be due to its free radical scavenging property (Lodhi et al., 2010). The anti allergic effect of *Tephrosia purpurea* has been reported (Gokhale and Saraf 2000). The extract of *Tephrosia purpurea* stabilized mast cells significantly showing its usefulness in the treatment and management of asthma (Gajera Paresch Lallubhai and Dalal Mittal, 2011). In another study *Tephrosia purpurea* showed spasmolytic activity in the trachea of guinea pigs thus strengthening the view of its use in asthma (Soni et al., 2004). *Tephrosia purpurea* has also been studied for its immunomodulatory effect (Damre et al., 2003). Ashokkumar et al., studied the diuretic activity of methanol extract of *Tephrosia purpurea* (Ashokkumar et al., 2012). The aqueous extract from roots of *Tephrosia purpurea* also posses antilithiatic activity

(Swathi et al., 2008). Still another study was made on the *Tephrosia purpurea* leaves for its protective and curative ability for renal injury in rats (Jain and Singhai, 2009). Study on chemical constituents of *Tephrosia candida* revealed a sesquiterpene having significant estrogenic activity (Hegazy et al., 2011). The chloroform and methanolic extract of *Tephrosia spinosa* showed significant ant helminthic activity against earth worms (*Pheretima posthuma*) (Ilango et al., 2011). The leaf extract of *Tephrosia vogelii* was found to possess significant anthelmintic activity against *Ascaridia galli*, a parasite in chicken (Siamba et al., 2007). The methanol extract of *Tephrosia vogelii* produced significant reduction in the blood pressure of cats (Audaudi et al., 2009).

## Conclusion and discussion

The plants of genus *Tephrosia* are of high therapeutic importance. We can see that a large number of species are studied for their chemical constituents but the number of isolated compounds from individual specie is very few with some exceptions. Mostly studied compounds include flavonoids and rotenoids. Studies on oil composition are very less. The genus has significant anticancer and larvicidal potential.

## References

- Wang YD, Tao MF, Wang L, Cheng WW, Wan XP. (2012). Selective regulation of osteoblastic OPG and RANKL by dehydroepiandrosterone through activation of the estrogen receptor  $\beta$ -mediated MAPK signaling pathway. *Hormone and Metabolic Research* 44, 494-500.
- Abreu PM, Luis MH. Constituents of *Tephrosia Uniflora*. (1996). *Natural Product Letters* 9, 81-86.
- Audaudi AO, Aluwong T, Salawu OA, Anuka J A. (2009). Blood pressure, analgesic and anti-inflammatory properties of methanolic extracts of *Tephrosia vogelii* in experimental animals. *Nigerian Veterinary Journal* 30, 37-43.
- Adinarayana K, Jayaveera KN, Madhu Katyayani B, Mallikarjuna Rao P. (2009). Growth inhibition and induction of apoptosis in estrogen receptor positive and negative human breast carcinoma cells by *Tephrosia calophylla* roots. *Pharmaceutical chemistry journal* 3, 35-41.
- Adinar-ayana K, Jayaveera KN, Rao PM, Chetty CM, Sandeep DK, Swetha C, Saleem TM. (2011). Acute toxicity and hepatoprotective effect of methanolic extract of *Tephrosia calophylla*. *Journal of medicinal plants research* 5, 266-273.
- Agarwal K, Gupta PK. (1983). Cytological studies in the genus *Tephrosia* Pers. *Cytologia* 48, 795-801.
- Ahmad A, Balakrishnan BR, Akhtar R, Pimprikar R. (2009). Antidiabetic activity of leaves of *Tephrosia villosa* Pers. in alloxan induced diabetic rats. *Journal of Pharmacy research* 2, 528-531.
- Ahmad S. (1986). Natural occurrence of *Tephrosia* flavones. *Phytochemistry* 25, 955-958.
- Ahmad VU, Ali Z, Hussaini SR, Iqbal F, Zahid M, Abbas M, Saba N. (1999). Flavonoids of *Tephrosia purpurea*. *Fitoterapia* 70, 443-445.
- Akande TO, Adeyeri MK, Longe OG. (2008). Nutritional and Economic Value of Leaf Meal (*Tephrosia bracteolata*) in Diet of Laying Hens. *Journal of Animal and Veterinary Advances* 7, 784-788.
- Akhtar R, Ahmad S, Khan M, Shaikh T, Mujawar T. (2011). Antihyperlipidemic effect of ethanolic extract of leaves of *Tephrosia purpurea* Linn in dexamethasone induced rats. [Article and details unavailable]
- Al-Zahrani R.M. (2007). Systematic Study of the Genus *Tephrosia* Pers. (Fabaceae) in Saudi Arabia. Post graduate thesis.
- Ammar NM, Jarvis BB. (1986). Major flavonoids of *Tephrosia nubica*. *Journal of Natural Products* 49, 719-720.

- Andrei CC, Ferreira DT, Faccione M, de Moraes LAB, de Carvalho MG, Braz-Filho R. (2000). C-prenylflavonoids from roots of *Tephrosia tunicata*. *Phytochemistry* 55, 799-804.
- Andrei CC, Vieira PC, Fernandes JB, da Silva MDG, Fo ER. (2002). New spirorotenoids from *Tephrosia candida*. *Zeitschrift. Fur Naturforschung C* 57, 418-422.
- Andrei CC, Vieira PC, Fernandes JB, Da Silva MFGF. (1997). Dimethylchromene rotenoids from *Tephrosia candida*. *Phytochemistry* 46, 1081-1085.
- Annalakshmi C, Satyabrata M, Suchandra G. (2009). The potential of *Tephrosia purpurea* as anti-*Helicobacter pylori* agent. *Journal of Ethnopharmacology* 124, 642-645.
- Arriaga AM, Lima JQ, Vasconcelos JN, de Oliveira MC, Lemos TL, Fonseca AM, Braz-Filho R. (2009). Antioxidant and larvicidal activities of *Tephrosia egregia* Sandw against *Aedes aegypti*. *Natural Product Communications* 4, 529-530.
- Arriaga AM, Magalhães FE, Feitosa EM, Malcher GT, Andrade-Neto M, Nascimento RF. (2005). Composition of the essential oil of *Tephrosia egregia* Sandw. *Journal of Essential Oil Research* 17, 451-452.
- Arriaga AMC, Lima JQ, Vasconcelos JN, de Oliveira MCF, Andrade Neto M, Santiago GMP, Braz-Filho R. (2009). Unequivocal assignments of flavonoids from *Tephrosia* sp.(Fabaceae). *Magnetic Resonance in Chemistry* 47, 537-540.
- Arriaga AMC, Malcher GT, Lima JQ, Magalhães FE, Gomes TM, Da Conceição M, Santiago GMP. (2008). Composition and Larvicidal Activity of the Essential Oil from *Tephrosia cinerea* Pers. *Journal of Essential Oil Research* 20, 450-451.
- Ashokkumar D, Narayana TV, Mazumder UK, Gupta M. (2012). Exploration of Diuretic Potential and Electrolyte Excretion of *Tephrosia purpurea* (Fabaceae) in Rats. *Journal of Dietary Supplements* 9, 9-18.
- Asuntha G, Prasannaraju Y, Sujitha D, Prasad KVSRRG. (2010). Assessment of effect of ethanolic extract of *Tephrosia purpurea* (Linn) Pers: fabaceae, activity on lithium-pilocarpine induced status epilepticus and oxidative stress in Wistar rats. *Brazilian journal of pharmacology* 2410.
- Atchison E. (1951). Studies in the Leguminosae. VI. Chromosome Numbers Among Tropical Woody Species. *American Journal of Botany* 38, 538-546.
- Babayemi J, Bamikole M. (2006). Supplementary value of *Tephrosia bracteolata*, *Tephrosia candida*, *Leucaena leucocephala* and *Gliricidia sepium* hay for West African dwarf goats kept on range. *Journal of Central European Agriculture* 7, 323-328.
- Babayemi OJ, Bamikole MA, Daniel IO, Ogungbesan A, Babatunde A. (2003). Growth, nutritive value and dry matter degradability of three *Tephrosia* species. *Nigerian Journal of Animal Production* 30, 62-70.
- Babayemi OJ, Bamikole MA. (2006). Effects of *Tephrosia candida* DC leaf and its mixtures with Guinea grass on in vitro fermentation changes as feed for ruminants in Nigeria. *Pakistan Journal of Nutrition* 5, 14-18.
- Babayemi OJ, Bamikole MA. (2007). Nutritive value of *Tephrosia candida* seed in West African dwarf goats. *Journal of Central European Agriculture* 7, 731-738.
- Balakrishnan BR, Sangameswaran B, Ahmed S, Bhaskar VH. (2007). Anti-hyperglycemic activity of roots of *Tephrosia villosa* Pers. *Plant Archives* 7, 729-731.
- Basu SP, Mandal JK, Mehdi NS. (1994). Anticonvulsant effect of pongamol. *Indian journal of pharmaceutical sciences* 56, 163-167.
- Bentley MD, Hassanali A, Lwande W, Njoroge PEW, Sitayo EN, Yatagai M. (1987). Insect antifeedants from *Tephrosia elata* deflers. *International Journal of Tropical Insect Science* 8, 85-88.
- BNLD Rangama, CL Abayasekara, GJ Panagoda, MRDM Senanayake. (2009). Antimicrobial activity of *Tephrosia purpurea* (Linn.) Pers. and *Mimosa elengi* (Linn.) against some clinical bacterial isolates. *J.Natn.Sci.Foundation Sri Lanka* 37, 139-145.
- Cai XA, Lai ZQ, Wang SC, Jiang JS, Yi XF, Wei JI, Liang YL. (2010). Study on Subchronic Toxicity Test of *Tephrosia candida* DC. on Mice. *China Animal Husbandry & Veterinary Medicine* 10, 11-15.



- Calixto JB, Santos AR, Yunes RA. (1998). A review of the plants of the genus *Phyllanthus*: their chemistry, pharmacology, and therapeutic potential. *Medicinal Research Reviews* 18, 225-258.
- Chakradhar V, Babu YH, Ganapaty S, Prasad YR, Rao NK. (2005). Anti-inflammatory activity of a flavonol glycoside from *Tephrosia spinosa*. *Natural Product Sciences* 11, 63-66.
- Chang LC, Chávez D, Song LL, Farnsworth NR, Pezzuto JM, Kinghorn AD. (2000). Absolute configuration of novel bioactive flavonoids from *Tephrosia purpurea*. *Organic letters* 2, 515-518.
- Chang LC, Gerhäuser C, Song L, Farnsworth NR, Pezzuto JM, Kinghorn AD. (1997). Activity-guided isolation of constituents of *Tephrosia purpurea* with the potential to induce the phase II enzyme, quinone reductase. *Journal of natural products* 60, 869-873.
- Chinniah A, Mohapatra S, Goswami S, Mahapatra A, Kar SK, Mallavadhani UV, Das PK. (2009). On the potential of *Tephrosia purpurea* as anti-*Helicobacter pylori* agent. *Journal of ethnopharmacology* 124, 642-645.
- Choudhary GP. (2007). In vitro antioxidant studies of the ethanolic extract of *Tephrosia purpurea* L. *Ancient Science of Life* 27, 26.
- Clark EP. (1930). Toxicarol a constituent of the South American fish poison cracca *Tephrosia Toxicaria*. *Journal of the American Chemical Society* 52, 2461-2464.
- Dagne E, Dinku B, Gray AI, Waterman PG. (1988). Pumilaisoflavones A and B from the seed pods of *Tephrosia pumila*. *Phytochemistry* 27, 1503-1505.
- Dagne E, Mammo W, Sterner O. (1992). Flavonoids of *Tephrosia polyphylla*. *Phytochemistry* 31, 3662-3663.
- Dagne E, Yenesew A, I Gray A, G Waterman P. (2012). Praecansone A: Evidence for the existence of 8, 9-(E) and 8, 9-(Z) isomers in extracts from *tephrosia pumila*. *Bulletin of the Chemical Society of Ethiopia* 4, 141-145.
- Dagne E, Yenesew A, Waterman PG. (1989). Flavonoids and isoflavonoids from *Tephrosia fulvivenis* and *Tephrosia pentaphylla*. *Phytochemistry* 28, 3207-3210.
- Damre AS, Gokhale AB, Phadke AS, Kulkarni KR, Saraf MN. (2003). Studies on the immunomodulatory activity of flavonoidal fraction of *Tephrosia purpurea*. *Fitoterapia* 74, 257-261.
- David Krupadanam GL, Sarma PN, Srimannarayana G, Subba Rao NV. (1977). New C-6 oxygenated rotenoids from *Tephrosia villosa* -- villosin, villosone, villol and villinol *Tetrahedron Letters* 18, 2125-2128.
- Deepak Kumar, P Dhamodaran, Nilani P, N Balakrishnan. (2012). Larvicidal activity of *Tephrosia purpurea*, (L) against the Larvae of *Culex quinquefasciatus*. *Journal of Applied Pharmaceutical Science* 2, 219-221.
- Delfel NE, Tallent WH, Carlson DG, Wolff IA. (1970). Distribution of rotenone and deguelin in *Tephrosia vogelii* and separation of rotenoid-rich fractions. *Journal of Agricultural and Food Chemistry* 18, 385-390.
- Deshpande SS, Shah GB. (2008). Pharmacological activity of *Tephrosia purpurea*. *American Association of Pharmaceutical Scientists Journal* 10, 10-15.
- Diwan R, Saxena R. (2010). Insecticidal property of flavinoid isolated from *Tephrosia purpuria*. *Int. J. Chem. Sci* 8, 777-782.
- Dominguez XA, Tellez O, Ramirez EG. (1983). Mixtecacin, a prenylated flavanone and oaxacacin its chalcone from the roots of *Tephrosia woodii*. *Phytochemistry* 22, 2047-2049.
- Duraipandiyan V, Ayyanar M, Ignacimuthu S. (2006). Antimicrobial activity of some ethnomedicinal plants used by Paliyar tribe from Tamil Nadu, India. *BMC complementary and alternative medicine* 6, 35.
- Dutt SK, Chibber SS. (1983). Candidol, a flavonol from *Tephrosia candida*. *Phytochemistry* 22, 325-326.
- Dzenda T, Ayo JO, Adelaiye AB, Auda AO, Ibrahim ND. (2008). Preliminary Investigation Into the acute oral toxicity of *Tephrosia vogelii* Leaves In Mice. *Nigerian Veterinary Journal* 28, 47-52.

- Dzenda T, Ayo JO, Adelaiye AB, Auda AO. (2008). Contraction of Isolated Guinea Pig Ileum By *Tephrosia vogelii* extract. *Nigerian Veterinary Journal* 27, 46-52.
- Dzenda T, Ayo JO, Adelaiye AB, Auda AO. (2007). Ethnomedical and veterinary uses of *Tephrosia vogelii* Hook F (Fabaceae): a review. *Niger Vet J* 28, 24-39.
- Falak AH, Shoeb A. (1987). A new epoxyflavanone from *Tephrosia hamiltonii*. *Planta Medica* 53, 220-221.
- Farnsworth NR, Akerele O, Bingel AS, Soejarto DD, Guo Z. (1985). Medicinal plants in therapy. *Bulletin of the world health organization* 63, 965-981.
- Gadzirayi C, Mutandwa E, Mwale M, Chindundu T. (2010). Utilization of *Tephrosia vogelii* in controlling ticks in dairy cows by small- scale commercial farmers in Zimbabwe. *African Journal of Biotechnology* 8, 4134-4136
- Gahlot K, Lal VK, Jha S. (2012). Phytochemical and Pharmacological potential of *Flemingia* Roxb. ex WT Aiton (Fabaceae). *International Journal of Phytomedicine* 3, 294-307.
- Gajera Pareshe Lallubhai, Dalal Mittal V. (2011). Mast cell stabilizing potential of the ethanolic extract of *Tephrosia purpurea* (LINN.) in the management of asthma. *IJRAP* 2, 1308-1312.
- Ganapaty S, Lakshmi P, Bobbarala V. (2010). Chemical and biological examination of the leaves of *Tephrosia tinctoria* PERS. *International Journal of Chemical and Analytical Science* 1, 14-17
- Ganapaty S, Lakshminarayana K, Lakshmi P, Thomas PS. (2009). *Asian Journal of Chemistry* 21, 1007-1010.
- Ganapaty S, Nyamathulla S, Srilakshmi GVK, Prasad KVNMR. (2008). Chemical and Antimicrobial Studies of the Roots of *Tephrosia villosa* (L) Pers. *Asian Journal of Chemistry* 20, 4498-4502.
- Ganapaty S, Pannakal ST, Srilakshmi GV, Lakshmi P, Waterman PG, Brun R. (2008). Pumilanol, an antiprotozoal isoflavanol from *Tephrosia pumil*. *Phytochemistry Letters* 1, 175-178.
- Ganapaty S, Srilakshmi GVK, Pannakal ST, Rahman H, Laatsch H, Brun R. (2009). Cytotoxic benzil and coumestan derivatives from *Tephrosia calophylla*. *Phytochemistry* 70, 95-99.
- Ganapaty S, Srilakshmi GVK, Thomas PS, Rajeswari NR, Ramakrishna S. (2009). Cytotoxicity and antiprotozoal activity of flavonoids from three *Tephrosia* species. *Journal of Natural Remedies* 9, 202-208.
- Geetha T, Varalakshmi P. (2001). Anti-inflammatory activity of lupeol and lupeol linoleate in rats. *Journal of ethnopharmacology* 76, 77-80.
- Gills LS. (1992). Ethnomedicinal uses of plants in Nigeria. Macmillan press Ltd, pp 229.
- Gokhale A, Saraf MN. (2000). Influence of ethanolic extract of *Tephrosia purpurea* Linn. on the late-phase of allergic reaction. *Indian Journal of Pharmaceutical Sciences* 62, 356-359.
- Gómez F, Calderón JS, Quijano L, Domínguez M, Ríos T. (1985). Viridiflorin, an isoflavone from *Tephrosia viridiflora*. *Phytochemistry* 24, 1126-1128.
- Gómez F, Quijano L, Calderón JS, Rodríguez C, Ríos T. (1985). Prenylflavans from *Tephrosia watsoniana*. *Phytochemistry* 24, 1057-1059.
- Gómez F, Quijano L, Garcia G, Calderón JS, Rios T. (1983). A prenylated flavan from *Tephrosia madrensis*. *Phytochemistry* 22, 1305-1306.
- Gómez-Garibay F, Calderón J, Quijano L, Téllez O, del Socorro Olivares M, Ríos T. (1997). An unusual prenyl biflavanol from *Tephrosia tepicana*. *Phytochemistry* 46, 1285-1287.
- Gómez-Garibay F, Calderón JS, De La O Arciniega M, Céspedes CL, Téllez-Valdés O, Taboada J. (1999). An unusual isopropenyldihydrofuran biflavanol from *Tephrosia crassifolia*. *Phytochemistry* 52, 1159-1163.
- Gómez-Garibay F, Quijano L, Calderón JS, Aguirre G, Ríos T. (1986). Abbotin and tephrobin, two new flavonoids from *Tephrosia abbotiae*. *Chemistry and Industry* 23, 827-832.
- Gómez-Garibay F, Quijano L, Calderón JS, Morales S, Rios T. (1988). Prenylflavanols from *Tephrosia quercetorum*. *Phytochemistry* 27, 2971-2973.
- Gómez-Garibay F, Quijano L, Hernandez C, Rios T. (1992). Enantiomultijugin, a flavone from *Tephrosia viciodes*. *Phytochemistry* 31, 2925-2926.

- Gomez-Garibay Federico, Tellez-Valdez Oswaldo, Moreno-Torres Gregorio, Calderon Josh S. (2002). Flavonoids from *Tephrosia major*. A new prenyl- $\beta$ -hydroxychalcone. *Zeitschrift für Naturforschung. C. A Journal of biosciences* 57, 579-583.
- Gunjegaonkar SM, Saraswathi CD, Hrishikeshavan HJ, Harish MS, Nargund LVG. (2010). Hepatoprotective and antioxidant activity of *Tephrosia purpurea* whole plant aqueous extract. *Pharmacologyonline* 2, 568-574.
- Gupta M, Mazumder UK, Gomathi P, Selvan V T. (2008). Antimicrobial activity of methanol extracts of *Plumeria acuminata* Ait. leaves and *Tephrosia purpurea* (Linn.) Pers. roots. *Natural Product Radianc* 7, 102-105.
- H Matovu and D. Olila. (2007). Larvicidal activity of *Tephrosia vogelii* crude extracts on mosquito larval stages. *Res.J.Biol.Sci* 2, 612-616.
- Hacker, J.B. (1990). A guide to herbaceous and shrub legumes of Queensland. University of Queensland Press, pp 275.
- Hari Kishore P, Vijaya Bhaskar Reddy M, Gunasekar D, Marthanda Murthy M, Caux C, Bodo B. (2003). A new coumestan from *Tephrosia calophylla*. *Chemical and pharmaceutical bulletin* 51, 194-196.
- Hegazy MEF, Abd El-Razek MH, Nagashima F, Asakawa Y, Paré PW. (2009). Rare prenylated flavonoids from *Tephrosia purpurea*. *Phytochemistry* 70, 1474-1477.
- Hegazy MEF, Mohamed AEHH, El-Halawany AM, Djemgou PC, Shahat AA, Paré PW. (2011). Estrogenic activity of Chemical Constituents from *Tephrosia candida*. *Journal of natural products* 74, 937-942.
- Hisham A, John S, Al-Shuaily W, Asai T, Fujimoto Y. (2006). (+)-Apollineanin: a new flavanone from *Tephrosia apollinea*. *Natural Product Research* 20, 1046-1052.
- Hu LZ, Li XH, Ma YQ, Yu X. (2011). Study on bacteriostatic activity of extracts in different solvents from *Tephrosia vogelii* Hook f. seeds. *Science and Technology of Food Industry* 1, 13-18.
- Hussain T, Siddiqui HH, Fareed S, Vijayakumar M, Rao CV. (2012). Chemopreventive evaluation of *Tephrosia purpurea* against N-nitrosodiethylamine-induced hepatocarcinogenesis in Wistar rats. *Journal of Pharmacy and Pharmacology* 64, 1195-1205.
- Ilango K, Xavier AR, Subburaju T. (2011). Anthelmintic Activity of Extracts of Aerial Parts of *Tephrosia spinosa* (Lf) Pres. *International Journal of Health Research* 4, 53-56.
- Ingham JL, Markham KR. (1980). Tephrocarpin, a pterocarpan phytoalexin from *Tephrosia bidwilli* and a structure proposal for acanthocarpan. *Phytochemistry* 21, 2969-2972.
- Irvine JE, Freyre RH. (1959). Source Materials for Rotenone, Occurrence of Rotenoids in Some Species of the Genus *Tephrosia*. *Journal of Agricultural and Food Chemistry* 7, 106-107.
- Jain A, Lodhi S, Singhai AK. (2009). Simultaneous estimation of quercetin and rutin in *Tephrosia purpurea* Pers by high performance thin-layer chromatography. *Asian Journal of Traditional Medicine* 4, 104- 109.
- Jain A, Singhai AK. (2009). Effect of *Tephrosia Purpurea* Pers on Gentamicin Model of Acute Renal Failure. 13th International Conference on Biomedical Engineering. Springer Berlin Heidelberg, pp 1438-1442.
- Jang DS, Park EJ, Kang YH, Hawthorne ME, Vigo JS, Graham JG, Kinghorn AD. (2003). Potential cancer chemopreventive flavonoids from the stems of *Tephrosia toxicaria*. *Journal of natural products* 66, 1166-1170.
- Jayaraman I, Ghanim A, Khan HA. (1980). A new prenylated flavanone [7-methylglabranin] from *Tephrosia villosa*. *Phytochemistry* 19.
- Juma WP, Akala HM, Eyase FL, Muiva LM, Heydenreich M, Okalebo FA, Yenesew A. (2011). Terpurinflavone: An antiplasmodial flavone from the stem of *Tephrosia Purpurea*. *Phytochemistry Letters* 4, 176-178.
- Kalume MK, Losson B, Angenot L, Tits M, Wauters JN, Frédéricich M, Saegerman C. (2012). Rotenoid content and in vitro acaricidal activity of *Tephrosia vogelii* leaf extract on the tick *Rhipicephalus appendiculatus*. *Veterinary Parasitology* 190, 204-209.

- Kannan RR, Vincent SGP. (2012). Rapid neurobehavioural analysis based on the effects of an acetylcholinesterase inhibitor from *Tephrosia purpurea* in Zebrafish. *Annals of Neurosciences* 19.
- Kare M, Kone MEK, Boulanger A, Niassy B, Lenouen D, Muckensturm B, Nongonierma A. (2006). Isolation, identification and antibacterial tests of chalcones and rotenoids of *Tephrosia deflexa* Baker. *J. Soc. ouest-afr. Chim* 11, 41-45.
- Khalafalah AK, Yousef AH, Esmail AM, Abdelrazik MH, Hegazy ME, Mohamed AEHH. (2010). Chemical constituents of *Tephrosia purpurea*. *Pharmacognosy research* 2, 72.
- Khalafallah AK, Suleiman SA, Yousef AH, El-kanzi NA, Mohamed AEHH. (2009). Prenylated flavonoids from *Tephrosia apollinea*. *Chinese Chemical Letters* 20, 1465-1468.
- Khalid SA, Waterman PG. (1981). 8- C -Prenylflavonoids from the seed of *Tephrosia bracteolata*. *Phytochemistry* 20, 1719-1720.
- Khalivulla SI, Reddy BAK, Gunasekar D, Blond A, Bodo B, Murthy MM, Rao TP. (2008). A new di-Oprenylated isoflavone from *Tephrosia tinctoria*. *Journal of Asian natural products research* 10, 953-955.
- Khan H, Chandrasek Haran K, Ghanim A. (1984). *Tephrosia falciformis*-a new source of triacontanol. *Ann. Arid Zone* 23, 39-40.
- Khan HA, Chandrasekharan I, Ghanim A. (1986). Falciformin, a flavanone from pods of *Tephrosia falciformis*. *Phytochemistry* 25, 767-768.
- Kim EM, Jung HR, Min TJ. (2001). Purification, structure determination and biological activities of 20 (29)-lupen-3-one from *Daedaleopsis tricolor* (Bull. ex Fr.) Bond. et Sing. *Bulletin-Korean Chemical Society* 22, 59-62.
- Kishore K, Roy D. (2011). *Tephrosia purpurea* Pers.(Fabaceae)-A common winter weed of Chhattisgarh, India-As a source of anticancer drug. *Indian J. Applied & Pure Bio* 26, 53-55.
- Koehn F, Carter GT. (2005). The evolving role of natural products in drug discovery. *Nature Reviews Drug Discovery* 4, 206-220.
- Kole RK, Satpathi C, Chowdhury A, Ghosh MR, Adityachaudhury N. (1992). Isolation of amorfolone, a potent rotenoid insecticide from *Tephrosia candida*. *Journal of Agricultural and Food Chemistry* 40, 1208-1210.
- Kumar AS, Amudha P, Kannan CS. (2011). Evaluation of anxiolytic activity of hydroalcoholic activity of *Tephrosia purpuria* (L) PERS on swiss albino mice. *International Journal of Pharmaceutical Sciences and research* 2, 1262-1269.
- Li XH, Huang XL, YU X, Zhu XY, Huang XH. (2010). Study on Scavenging Free Radical Activity of Extracts from *Tephrosia vogelii* Hook f. Seeds. *Food Research and Development* 1, 5-9.
- Liu J. (1995). Pharmacology of oleanolic acid and ursolic acid. *Journal of ethnopharmacology* 49, 57-68.
- Lodhi S, Pawar RS, Jain AP, Jain A, Singhai AK. (2010). Effect of *Tephrosia purpurea* (L) Pers. on Partial Thickness and Full Thickness Burn Wounds in Rats. *Journal of Complementary and Integrative Medicine* 7, 1-15.
- Lwande W, Bentley MD, Macfoy C, Lugemwa FN, Hassanali A, Nyandat E. (1987). A new pterocarpan from the roots of *Tephrosia hildebrandtii*. *Phytochemistry* 26, 2425-2426.
- Lwande W, Greene CS, Bentley MD. (1985). Flavonoids from the roots of *Tephrosia elata*. *Journal of Natural Products* 48, 1004-1005.
- Lwande W, Hassanali A, Bentley MD, Monache FD. (1986). 8-C-Prenylated Flavones from the Roots of *Tephrosia hildebrandtii*. *Journal of Natural Products* 49, 1157-1158.
- Lwande W, Hassanali A, Njoroge PW, Bentley MD, Delle Monache F, Jondiko JI. (1985). A new 6 ahydroxypterocarpan with insect antifeedant and antifungal properties from the roots of *Tephrosia hildebrandtii* Vatke. *Insect science and its application* 6, 537-541.
- Machocho AK, Lwande W, Jondikoa JI, Moreka LVC, Hassanali A. (1995). Three new flavonoids from the root of *Tephrosia emoroides* and their antifeedant activity against the larvae of the spotted stalk borer *Chilo partellus* Swinhoe. *Pharmaceutical Biology* 33, 222-227.

- Madhusudhana J, Reddy RN, Reddy BAK, Reddy MVB, Gunasekar D, Deville A, Bodo B. (2010). Two new geranyl flavanones from *Tephrosia villosa*. *Natural Product Research* 24, 743-749.
- Maldini M, Montoro P, Macchia M, Pizza C, Piacente S. (2011). Profiling of Phenolics from *Tephrosia cinerea*. *Planta medica* 77, 1861-1864.
- Martin JT. (1936). Occurrence of rotenone in *Tephrosia macropoda* Harv. *Nature* 137, 1075-1080.
- Martinez RM, Zarpelon AC, Zimmermann VV, Georgetti SR, Baracat MM, Fonseca MJ, Casagrande R. (2012). *Tephrosia sinapou* extract reduces inflammatory leukocyte recruitment in mice: effect on oxidative stress, nitric oxide and cytokine production. *Revista Brasileira de Farmacognosia* 22, 587-597.
- Mbomi SE, Ogungbesan AM, Babayemi OJ, Nchinda VP. (2011). Chemical composition, acceptability of three *Tephrosia* species and use of *Tephrosia purpurea* as supplement for grazing animals in the western highlands of Cameroon. *Journal of Environmental Issues and Agriculture in Developing Countries* 3, 132-139.
- Mohan C.U. (2011). Antihyperlipidemic effect of *Tephrosia calophylla* against acute and chronic hyperlipidemia in wistar albino rats. Post graduate thesis.
- Monache FD, Labbiento L, Marta M, Lwandet W. (1986). 4 $\beta$ -substituted flavans from *Tephrosia hildebrandtii*. *Phytochemistry* 25, 1711-1713.
- Muiva LM, Yenesew A, Derese S, Heydenreich M, Peter MG, Akala HM, Walsh D. (2009). Antiplasmodial  $\beta$ -hydroxydihydrochalcone from seedpods of *Tephrosia elata*. *Phytochemistry Letters* 2, 99-102.
- Muiva LM. (2012). Antiplasmodial and Larvicidal Flavonoids from the Seedpods of *Tephrosia elata* and *Tephrosia aequilata*. Postgraduate thesis.
- Murthy MSR, Rao EV. (1985). Maxima isoflavone J: a new O-prenylated isoflavone from *Tephrosia maxima*. *Journal of natural products* 48, 967-968.
- Niassy B, Um BH, Lobstein A, Weniger B, Koné M, Anton R. (2004). Flavonoïdes de *Tephrosia deflexa* et *Tephrosia albifoliolis*. *Comptes Rendus Chimie* 7, 993-996.
- Nile SH, Khobragade CN. (2011). Indian Phytochemical analysis, antioxidant and xanthine oxidase inhibitory activity of *Tephrosia purpurea* Linn. root extract. *Journal of Natural Products and Resources* 2, 52-58.
- Nisar MF, Ismail S, Arshad M, Majeed A, Arfan M. (2011) Ethnomedicinal Flora of District Mandi Bahaudin, Pakistan. *Middle-East Journal of Scientific Research* 9, 233-238.
- Nondo RS, Mbwambo ZH, Kidukuli AW, Innocent EM, Mihale MJ, Erasto P, Moshi MJ. (2011). Larvicidal, antimicrobial and brine shrimp activities of extracts from *Cissampelos mucronata* and *Tephrosia villosa* from coast region, Tanzania. *BMC complementary and alternative medicine* 11, 33-37.
- Onaolapo MOA, Nzelibe HC, Aduadi AO, Ayo JO. (2009). Toxicity and antipyretic studies of the crude extract of *Tephrosia bractiolata* leaves. *Journal of Phytomedicine and Therapeutics*. 9,
- Parmar VS, Jain R, Gupta SR, Boll PM, Mikkelsen JM. (1988). Phytochemical Investigation of *Tephrosia candida*: Hplc Separation of Tephrosin and 12a-Hydroxyrotenone. *Journal of Natural Products* 51, 185-185.
- Parmar VS, Rathore JS, Jain R, Henderson DA, Malone JF. (1989). Occurrence of pongamol as the enol structure in *Tephrosia purpurea*. *Phytochemistry* 28, 591-593.
- Patel Anuradha Jitendra, Thakor Akanksha Pravin. (2012). Prospective use of *Tephrosia Purpurea* in Remedial Treatment of PCOS: Study in Wistar Rat. *ISCA J. Biological Sci* 1, 1-6.
- Pavana P, Manoharan S, Renju GL, Sethupathy S. (2007). Antihyperglycemic and antihyperlipidemic effects of *Tephrosia purpurea* leaf extract in streptozotocin induced diabetic rats. *Journal of Environmental Biology* 28, 833-837.
- Pavana P, Sethupathy S, Santha K, Manoharan S. (2009). Effects of *Tephrosia purpurea* aqueous seed extract on blood glucose and antioxidant enzyme activities in streptozotocin induced diabetic rats. *African Journal of Traditional, Complementary, and Alternative Medicines* 6, 78.

- Pelter A, Ward RS, Rao EV, Raju NR. (1981). 8-Substituted flavonoids and 3'-substituted 7-oxyg-enated chalcones from *Tephrosia purpurea*. *J. Chem. Soc., Perkin Trans 1*, 2491-2498.
- Pirrung MC, Lee YR. (1995). Total Synthesis and Absolute Configuration of Pseudosemiglabrin, a Platelet Aggregation Antagonist, and Its Diastereomer Semiglabrin. *Journal of the American Chemical Society* 117, 4814-4821.
- Prabhakar P, Vanangamudi A, Gandhidasan R, Raman PV. (1996). Hookerianin: a flavone from *Tephrosia hookeriana*. *Phytochemistry* 43, 315-316.
- Prashant A, Krupadanam GD. (1993). A new prenylated dehydrorotenoid from *Tephrosia villosa* seeds. *Journal of natural products* 56, 765-766.
- Prashant A, Krupadanam GL. (1993). Dehydro-6-hydroxyrotenoid and lupenone from *Tephrosia villosa*. *Phytochemistry* 32, 484-486.
- Quijano L, Rios T. (1991). Flavanones from *Tephrosia leiocarpa*. *Phytochemistry* 30, 3832-3834.
- Qureshi R, Bhatti GR, Memon RA. (2010). Ethnomedicinal uses of herbs from northern part of NARA desert, Pakistan. *Pak. J. Bot* 42, 839-851.
- Rajal Shah, Sachin Parmar, Punit Bhatt, Sumitra Chanda. (2011). Evaluation of hepatoprotective activity of ethyl acetate fraction of tephrosia purpurea. *Pharmacologyonline* 3, 188-194.
- Rajani P, Sarma N. (1988). A coumestone from the roots of *Tephrosia hamiltonii*. *Phytochemistry* 27, 648-649.
- Rangaswami S, Rao GR. (1955). Occurrence of rutin in the leaves of *Tephrosia lanceolata*. *Indian J. Pharm* 17, 136-137.
- Rao EV, Murthy M, Ward RS. (1984). Nine isoflavones from *Tephrosia maxima*. *Phytochemistry* 23, 1493-1501.
- Rao EV, Prasad YR. (1992). Flavonoids from *Tephrosia spinosa*. *Phytochemistry* 32, 183-185.
- Rao EV, Prasad YR. (1992). Two chalcones from *Tephrosia spinosa*. *Phytochemistry* 31, 2121-2122
- Rao EV, Raju NR. (1979). Occurrence of (-)-isolonchocarpin in the roots of *Tephrosia purpurea*. *Phytochemistry* 18, 1581-82.
- Rao EV, Raju NR. (1984). Two flavonoids from *Tephrosia purpurea*. *Phytochemistry* 23, 2339-2342.
- Rao PP, Srimanarayana G. (1981). Tephtrinone—a new flavone from *Tephrosia villosa*. *Curr. Sci* 50, 319-320.
- Ratsimamanga-Urverg S, Rasoanaivo P, Rabemanantsoa C, Rakoto Ratsimamanga A, Frappier F. (1994). Antibacterial activity of flavonoids isolated from *Mundulea monantha* and *Tephrosia linearis*. *Fitoterapia* 65, 551-553.
- Reddy RV, Khalivulla SI, Reddy BA, Reddy MV, Gunasekar D, Deville A, Bodo B. (2009). Flavonoids from *Tephrosia calophylla*. *Natural product communications* 4, 59-62.
- Rout S.P, Choudary KA, Kar DM, Das Lopamu dra, Jain Avijeet. (2009). Plants in traditional medicinal system-future source of new drugs. *Int. J. Pharm. Pharm. Sci* 1, 1-23.
- Roy M, Bhattacharya PK, Pal S, Chowdhuri A, Adityachaudhury N. (1987). Dehydrodihydrorotenone and flemichappar-B in *Tephrosia candida*. *Phytochemistry* 26, 2423-2424.
- Roy M, Mitra SR, Bhattacharyya A, Adityachaudhury N. (1986). Candidone, a flavanone from *Tephrosia candida*. *Phytochemistry* 25, 961-962.
- Sandhya S, Venkatramana K, Vinod KR, RK Chaitanya, J Chandrasekhar, K Sudhakar, T Rajeswar. (2010). Membrane stabilizing potency of two *Tephrosia* species. *J Phytol* 2, 42-46.
- Divya S, Haritha V, Prasad KVSRG. (2011). Evaluation of tephrosia calophylla for antiulcer activity in experimental rats. *Pharmacologyonline* 3, 573-585.
- Santos JCA, Riet-Correa F, Dantas AF, Barros SS, Molyneux RJ, Medeiros RM, de Oliveira OF. (2007). Toxic hepatopathy in sheep associated with the ingestion of the legume *Tephrosia cinerea*. *Journal of Veterinary Diagnostic Investigation* 19, 690-694.
- Saxena VK, Choubey A. (1997). A novel neoflavonoid glycoside from *Tephrosia purpurea* stem. *Fitoterapia* 68, 359-360.

- Shanmugapriya R, Umamaheswari G, Thirunavukkarasu P, Renugadevi G, Ramanathan T. (2011). Cytotoxic effect of *Tephrosia purpurea* extracts on HELA cervical cancerous cell line. *Inventi Rapid: Molecular Pharmacology* Vol. 2011, Article ID-"Inventi:mp/49/11"
- Sharaby A, Ammar N. (1997). Biological activity of extracts of *Tephrosia nubica* (Boiss) Baker against *Spodoptera littoralis* (Boisd.) and *Agrotis ipsilon* (Hufn.). *Tropenlandwirt* 98, 143-150.
- Sharma P, Rastogi S, Bhatnagar S, Srivastava J K, Dube A, Guru PY, Dhawan BN. (2003). Antileishmanial action of *Tephrosia purpurea* linn, extract and its fractions against experimental visceral leishmaniasis. *Drug development research* 60, 285-293.
- Sharma SK, Vasudeva N, rathi P. (2008). Isolation and identification of a new phytoester from *Tephrosia purpurea* (Linn.) Pers. Root. *Int. J. Chem. Sci* 6, 1734-1741.
- Sharma VM, Rao PS. (1992). A prenylated chalcone from the roots of *Tephrosia spinosa*. *Phytochemistry* 31, 2915-2916.
- Siamba DN, Okitoi LO, Watai MK, Wachira AM, Lukibisi FB, Mukisira EA. (2007). Efficacy of *Tephrosia vogelli* and *Vernonia amygdalina* as anthelmintics against *Ascaridia galli* in indigenous chicken. *Livestock Research for Rural Development* 19, 12.
- Sinha B, Natu AA, Nanavati DD. (1982). Prenylated flavonoids from *Tephrosia purpurea* seeds. *Phytochemistry* 21, 1468-1470.
- Smalberger TM, Van Den Berg AJ, Vleggaar R. (1973). Flavonoids from *Tephrosia*—VI: The structure of semiglabin and semiglabinol. *Tetrahedron* 29, 3099-3104.
- Smalberger TM, Vleggaar R, Weber JC. (1974). Flavonoids from *Tephrosia* —VII: The constitution and absolute configuration of lupinifolin and lupinifolinol, two flavanones from *Tephrosia lupinifolia* Burch (DC). *Tetrahedron* 30, 3927-3931.
- Smalberger TM, Vleggaar R, Weber JC. (1975). Flavonoids from *Tephrosia* —VIII: The structure of elongatin, an isoflavone from *Tephrosia elongata* E. Mey. *Tetrahedron* 31, 2297-2301.
- Soni KK, Khare ML, Saxena RC. (2004). Spasmolytic activity of a herbal drug isolated from *Tephrosia purpurea* in guinea pigs. *Ancient Science of Life* 23, 59-65.
- Subhadra S, Kanacharalapalli VR, Ravindran VK, Parre SK, Chintala S, Thatipally R. (2011). Comparative toxicity assessment of three *Tephrosia* species on *Artemia salina* and animal cell lines. *Journal of Natural Pharmaceuticals* 2, 143-148.
- Suliman HB, Wasfi IA, Adam SEI. (1982). The toxic effects of *Tephrosia apollinea* on goats. *Journal of Comparative Pathology* 92, 309-315.
- Swathi D, Sujatha D, Bharathi K, Prasad KVSRRG. (2008). Antilithiatic activity of the aqueous extract of the roots of *Tephrosia purpurea* Linn. *Pharmacognosy Magazine* 4, 206-211.
- Talib Hussain, Sheeba Fareed, HH Siddiqui, M Vijaykumar, Chandana Venkateswara Rao. (2012). Acute and subacute oral toxicity evaluation of *Tephrosia purpurea* extract in rodents. *Asian Pacific Journal of Tropical Disease* 129-132.
- Tarus PK, Machocho AK Chhabra SC. (2002). Flavonoids from *Tephrosia aequilata*. *Phytochemistry* 60, 375-379.
- Tatteksfield F, Gimingham CT. (1932). The insecticidal properties of *Tephrosia magropoda* HARV. And other tropical plants. *Annals of Applied Biology* 19, 253-262.
- Udeani GO, Gerhäuser C, Thomas CF, Moon R C, Kosmeder JW, Kinghorn AD, Pezzuto JM. (1997). Cancer chemopreventive activity mediated by deguelin, a naturally occurring rotenoid. *Cancer research* 57, 3424-3428.
- Valli G, Vasanthi A, Vijayalakshmi R, Thanga Thirupathi A. (2011). Antipyretic and anti-inflammatory activities of *Tephrosia purpurea* root extracts. *International Journal of pharmaceutical Research and development* 3, 211-217.
- Vanangamudi A, Gandhidasan R, Raman PV. (1997). Eupalitin 3-O-b-D-galactopyranoside from *Tephrosia spinosa*. *Fitoterapia* 68, 560-564.
- Vanangamudi A, Prabhakar P, Gandhidasan R, Raman PV. (1997). Tephrorianin: a novel flavone from *Tephrosia hookeriana*. *Fitoterapia* 68, 543-544.

- Vasconcelos JN, Lima JQ, Lemos TLGD, Oliveira MDCFD, Almeida MMB, Andrade-Neto M, Braz-Filho R. (2009). Chemical and biological study of the *Tephrosia toxicaria* Pers. *Química Nova* 32, 382-386.
- Venkata Rao E, Rajendra Prasad Y, Sree Rama Murthy M. (1994). A prenylated flavanone from *Tephrosia maxima*. *Phytochemistry* 37, 111-112.
- Venkata Rao E, Sree Rama Murthy M. (1985). Further studies on the isoflavones of *Tephrosia maxima*. *Phytochemistry* 24, 875-876.
- Venkata Rao E, Venkataratnam G, Vilain C. (1985). Flavonoids from *Tephrosia fulvinervis* *Phytochemistry* 24, 2427-2430.
- Venkataratnam G, Rao EV, Vilain C (1987). Flavonoids of *Tephrosia procumbens*—revised structure for praecansone A and conformation of praecansone. *BJ. Chem. Soc., Perkin Trans 1*, 2723-2727.
- Venkataratnam G, Rao EV, Vilain C. (1986). Fulvinervin C, a flavone from *Tephrosia fulvinervis*. *Phytochemistry* 25, 1507-1508.
- Vilain C. (1980). Barbigerone, a new pyranoisoflavone from seeds of *Tephrosia barbigeri*. *Phytochemistry* 19, 988-989.
- Vilain C. (1983). Isopongaflavone from *Tephrosia barbigeri* seeds. *Pharmazie* 38, 876-877.
- Virupanagouda P.Patil, Shivakumar Hugar, Nanjappaiah HM, Navanath Kalyane, Mohan Chowdhary, Pandarinath. (2011). Phytopharmacology of *Tephrosia purpurea* Linn: An Overview. *Pharmacologyonline* 3, 1112-1140.
- Vleggaar R, Kruger GJ, Smalberger TM, Van den Berg AJ. (1978). Flavonoids from *Tephrosia* —XI: The structure of glabratephrin. *Tetrahedron* 34, 1405-1408.
- Vleggaar R, Smalberger TM, Van den Berg AJ. (1975). Flavonoids from *Tephrosia*—IX: The structure of multijugin and multijuginol. *Tetrahedron* 31, 2571-2573.
- Wanga BN, Akenga T, Imbuga M, Gitonga L, Olubayo F, Namungu P. (2007). Antimicrobial activity of extracts from *Tephrosia vogelii* Hook F. *Journal of Agriculture, Science and Technology* 8, 1-14.
- Wanjala FME, Oriedo RA, Karanja DMS. (2006). The larvicidal efficacy of *Tephrosia nyikensis* Bak subsp. *victoriensis* Gellet and Brummit crude leaf extract on *Anopheles* mosquitoes. *Discovery and Innovation* 17, 56- 59.
- Waterman PG, Khalid SA. (1980). The major flavonoids of the seed of *Tephrosia apollinea*. *Phytochemistry* 19, 909-915.
- Wei HH, Xu HH, Xie HH, Xu LX, Wei XY. (2009). Sesquiterpenes and lignans from *Tephrosia vogelii*. *Helvetica Chimica Acta* 92, 370-374.
- Were O, Munavu RM, Lwande W, Nyandat E. (1990). Flavonoids from *Tephrosia interrupta* and *T. linearis*. *Fitoterapia* 61, 372.
- Willis, J. C. (1973). *The Dictionary of Flowering Plants and Ferns*. 8th ed. Cambridge University Press, Cambridge, UK, pp 1135.
- Yang D, Michel L, Chaumont JP, Millet-Clerc J. (2000). Use of caryophyllene oxide as an antifungal agent in an in vitro experimental model of onychomycosis. *Mycopathologia* 148, 79-82.
- Yenesew A, Dagne E, Waterman PG. (1989). Flavonoids from the seed pods of *Tephrosia pumila*. *Phytochemistry* 28, 1291-1292.
- Yu X, Yan Z, Li X, Huang X, Zeng X. (2009). Analysis of Physical Characteristics, Composition of *Tephrosia vogelii* Hook f. Seeds and its Oil Component [J]. *Acta Scientiarum Naturalium Universitatis Sunyatseni* 1, 11-17.